Journal of the Marine Biological Association of the United Kingdom, 2009, 89(1), 179–201. ©2008 Marine Biological Association of the United Kingdom
doi:10.1017/S0025315408002476 Printed in the United Kingdom
SC/N10/CC For Info 2

REVIEW

Potential impacts of climate change and greenhouse gas emissions on Mediterranean marine ecosystems and cetaceans

D.D. GAMBAIANI[1], P. MAYOL[1], S.J. ISAAC[2] AND M.P. SIMMONDS[3]

[1]Souffleurs d'Ecume, Mairie de la Celle, 83170 La Celle, France, [2]Halcrow, Burderop Park, Swindon, Wiltshire, SN4 0QD, [3]The Whale and Dolphin Conservation Society, Brookfield House, Chippenham, Wiltshire, SN15 1LJ, UK

The combustion of fossil fuels and the resultant impacts on climate may now represent one of the largest environmental threats. In the Mediterranean Sea, changes in bio-chemical and physical seawater properties resulting from global warming are likely to alter marine biodiversity and productivity, trigger trophic web mismatches and encourage diseases, toxic algal bloom and propagation of thermophilic species. This review highlights the current and potential threats of climate change to the Mediterranean marine ecosystems, including cetaceans, and stresses the emergent necessity for more integrated regulations and policies for the protection of marine biodiversity. For instance, in the Mediterranean Sea, the distribution and abundance of the small euphausid species Meganyctiphanes norvegica *is correlated with specific hydrobiological parameters including seawater temperature, salinity and current patterns. Situated at the northern limit of its ecological tolerance, this species, which constitutes the only known food supply of the fin whales* (Balaenoptera physalus) *in this region, might be affected by climate change-induced alteration of ocean circulation.*

Keywords: potential impacts, climate change, greenhouse gas emissions, Mediterranean, marine ecosystems, cetaceans

Submitted 31 July 2007; accepted 3 June 2008; first published online 23 September 2008

INTRODUCTION

Until recently, scientists have hesitated to associate the global warming phenomenon with anthropogenic greenhouse gas emissions. However, a strong correlation has been demonstrated along with other supporting evidence and it is now generally accepted that greenhouse gas emissions are responsible for global warming (e.g. Tett *et al.*, 2000; Smith *et al.*, 2005; IPCC, 2007a). We know that our carbon dioxide emissions, which represent ~7 gigatons (Gt) of carbon per year (Bopp *et al.*, 2002), are responsible for disastrous ecological disorders. Each year, 2 Gt of carbon is absorbed by the oceans through physical and biological processes but, this 'carbon pump' could be saturated by 2020–2030 (Orr *et al.*, 2001). A significant reduction in the efficiency of the Southern Ocean sink of CO_2 has recently been observed (Le Quéré *et al.*, 2007). Studies indicate that climate/carbon cycle feedbacks could be responsible for an additional increase of the partial pressure of CO_2 in the atmosphere of 10% to 50% (Cox *et al.*, 2000). Working Group 1 of the Intergovernmental on Climate Change (IPCC) shows that, according to different scenarios, the average warming of the planet will be between 1.8 and 4°C by 2100 (IPCC, 2007b). Such an increase of temperature in a very short time period represents a real 'climatic shock' for which no solution has yet been considered by our society (Jancovici, 2005).

Positive feedback processes would tend to accelerate the rate of global warming (Woodwell & Mackenzie, 1995). The amount of methane sequestered as methane hydrates in the oceanic sediments of continental plates, as well as in permafrost regions, could be 3000 times larger than the amount of methane present in the atmosphere (Blunier, 2000). With an increase of water temperatures, methane hydrates could become unstable and release vast quantities of methane into the atmosphere. Given that methane is 20 times more efficient than carbon dioxide at trapping infrared radiation, this could have catastrophic climatic consequences. Such an event, known as a 'methane burp', occurred about 200 million years ago. By increasing the atmospheric temperature by 4 to 8 degrees, and reducing the amount of oxygen, this phenomenon may have been responsible for the extinction of 80% of the oceanic fauna at that time (Hesselbo *et al.*, 2000; Kennett *et al.*, 2000).

Atmospheric carbon dioxide is also known to be absorbed by the oceans, leading to ocean acidification. This process is already

Corresponding author:
D.D. Gambaiani
Email: delphine.gambaiani@souffleursdecume.com

SC/N10/CC For Info 2

affecting surface waters (Royal Society, 2005) and future increases in atmospheric carbon dioxide via the combustion of fossil fuels are expected to profoundly affect ocean chemistry and marine life (Caldeira & Wickett, 2003; Feely *et al.*, 2004; Royal Society, 2005; Orr *et al.*, 2005; Bass *et al.*, 2006).

Global warming is already affecting marine ecosystems (CBD, 2003; Parmesan & Yohe, 2003). According to Thomas *et al.* (2004), climate change is today one of the most serious threats to biodiversity and 'on the basis of mid-range climate warming scenarios for 2050, 15–37% of species in our sample of regions and taxa will be "committed to extinction"'.

As a miniature ocean (0.82% of the world's ocean surface) with fast turnover time (40–50 years) the semi-enclosed Mediterranean, which accounts for 8% to 9% of global marine biodiversity and contains numerous endemic species (Bianchi & Morri, 2000), is likely to rapidly respond to external forcing like climate change (Béthoux & Gentili, 1996; Monaco, 1998; Béthoux *et al.*, 1999; Turley, 1999; Simmonds & Nunny, 2002; Giorgi, 2006).

The aim of this review is to illustrate the links existing between biodiversity and climate and demonstrate how anthropogenic greenhouse warming can affect the Mediterranean marine ecosystems including large predators such as cetaceans. It illustrates the impacts of global warming on abiotic and biotic systems in the Mediterranean Sea and stresses the necessity to consider climate change as a major issue.

DISCUSSION

Influence of climate change on Mediterranean abiotic parameters

By influencing climatic features (e.g. atmospheric temperature, precipitation budget and extreme meteorological events), global warming is likely to affect the chemical and physical properties of Mediterranean waters.

Present and future trends of Mediterranean climate and hydrology

At the beginning of the 20th Century, the temperature and salinity of the Western Mediterranean Deep Water (WMDW) was practically constant (Lacombe *et al.*, 1985; Béthoux *et al.*, 1990). Over the last few decades, numerous studies have indicated a correlation between climate patterns and changes in seawater properties, with general increases in both temperature and salinity over the entire Mediterranean (Lacombe *et al.*, 1985; Béthoux *et al.*, 1990; Leaman & Schott, 1991; Rohling & Bryden, 1992; Francour *et al.*, 1994; Sparnocchia *et al.*, 1994; Astraldi *et al.*, 1995; Graham, 1995; Béthoux & Gentili, 1996; Béthoux *et al.*, 1998, Krahmann & Schott, 1998; Duarte *et al.*, 1999; Danovaro *et al.*, 2001; Astraldi *et al.*, 2002; Fuda & Millot, 2002; Gertman & Hecht, 2002; Goffart *et al.*, 2002; Lascaratos *et al.*, 2002; Lopez-Jurado, 2002; Manca *et al.*, 2002; Prieur, 2002; Salat & Pascual, 2002; Vargas-Yanes *et al.*, 2002; Vilibic, 2002; Walther *et al.*, 2002; Rixen *et al.*, 2005; Li *et al.*, 2006; Millot *et al.*, 2006; Somot *et al.*, 2006). It is now assumed that the WMDW has become warmer and saltier in response to anthropogenic greenhouse effects and changes in the freshwater budget (Béthoux *et al.*, 1990; Rohling & Bryden, 1992; Graham, 1995; Béthoux *et al.*, 1998).

Furthermore, from 1900 to 2000, the Mediterranean experienced a 10% decrease in summer rainfall (IPCC, 2001a). These trends are expected to persist and intensify over time, and extreme meteorological events such as droughts (e.g. in the summers of 1995 and 2003) and flooding will become more frequent in the future (e.g. Parry, 2000; IPCC, 2001b; Li, 2003; EEA, 2004; Giannakopoulos *et al.*, 2005; Xoplaki *et al.*, 2005; Luterbacher *et al.*, 2007). By the 2020s, under high CO_2 scenarios, summer in southern Europe will be as hot as or hotter than the 2003 summer (Luterbacher *et al.*, 2004).

Font *et al.* (2007) have documented that the exceptional winter of 2005 affected the deep-water thermohaline properties of the north-western Mediterranean. Similarly, in the Aegean Sea, it is likely that changes in the freshwater budget could have modified seawater salinity and altered water-mass circulation (Roether *et al.*, 1996; Theocharis *et al.*, 1999). Climate variation—including extreme events—is not necessarily a result of directional climate change. However, climate change is likely to create more extreme meteorological events (e.g. storms and drought), and these can alter the biochemical and physical properties of seawater (Poumadère *et al.*, 2005; Olita *et al.*, 2007).

SEA LEVEL RISE

Ecologically and socio-economically rich coastal marine systems are under threat from anthropogenic global warming (IPCC, 2001a; Roessig *et al.*, 2004). Sea level rise resulting from climate change-related processes such as seawater thermal expansion or glacier melt will trigger coastal flooding (Paskoff, 2001). According to Tsimplis & Rixen (2002) from 1993 onwards, the warming of surface waters in the Eastern Mediterranean has caused sea level rise. In the last century, sea level has risen by 0.10 to 0.20 m around Europe (IPCC, 2001a) and an increase of sea level of 3.3 cm in 11 years (0.3 cm per year) was recorded at the oceanographic and meteorological station at l'Estartit in north-east Spain (Salat & Pascual, 2002). Between 1990 and 2100, the predicted trend for sea level rise is 2.2 to 4.4 times higher than the rate recorded in the 20th Century (IPCC, 2001a). Consequently, some Mediterranean coastal areas and wetlands, such as the Camargue, are threatened (Nicholls & Hoozemans, 1996; Pfeifle *et al.*, 2004).

ATMOSPHERIC CHANGE AND CLIMATE IMPACTS IN THE MEDITERRANEAN

Anthropogenic climate change tends to influence large atmospheric patterns like the North Atlantic Oscillation (NAO) and the El Niño Southern Oscillation (ENSO) (Timmermann *et al.*, 1999; Visbeck *et al.*, 2001; Würsig *et al.*, 2002; Cohen & Barlow, 2005). Global warming is likely to contribute to the positive phase of the NAO, which was responsible for warmer winters in Europe in the last 10–20 years (Hurrell, 1996; Visbeck *et al.*, 2001; Gillet *et al.*, 2003; Cohen & Barlow, 2005) and El Niño conditions may be more frequent with global warming (Timmermann *et al.*, 1999; Würsig *et al.*, 2002).

The Mediterranean climate is influenced by large-scale atmospheric circulation systems including the inter-decadal NAO and ENSO (Hurrell, 1995; Raicich *et al.*, 2001; Mariotti *et al.*, 2002; Lionello & Sanna, 2005). Changes in Mediterranean seawater temperature and salinity have been associated with the NAO (e.g. Béthoux & Gentili, 1999; Béthoux *et al.*, 1999; Tsimplis *et al.*, 2006). In fact, the

SC/N10/CC For Info 2

NAO influences Mediterranean precipitation and river discharges (nutrient inputs), which tend to be lower during positive NAO episodes (Lloret *et al.*, 2001; Struglia *et al.*, 2004; Xoplaki *et al.*, 2004, 2006). According to Tsimplis & Josey (2001), NAO is also influencing sea level change in the Mediterranean Sea. Mariotti *et al.* (2002) have also identified a link between ENSO and Mediterranean autumn rainfall anomalies. The relationship between NAO, ENSO and climate change is an important issue and requires further study.

Climate change impacts on Mediterranean marine biodiversity and ecosystems

Evidence has demonstrated that climate variation and changes in the properties of ecological systems are strongly correlated (e.g. Orr *et al.*, 1992; Astraldi *et al.*, 1995; Bombace, 2001; Hannah *et al.*, 2002; Walther *et al.*, 2002; Duffy, 2003; Root *et al.*, 2003; Roessig *et al.*, 2004) and that global warming is already affecting numerous marine species throughout the world (Table 1). In the Mediterranean, it is now acknowledged that climate change-induced temperature variations have altered biological patterns and biodiversity (e.g. Francour *et al.*, 1994; Turley, 1999; Bianchi & Morri, 2000).

Table 1. Examples of species affected by climate change induced-factors around the world (Mediterranean Sea excluded).

Areas	Examples of affected species
North-east Atlantic	• Marine flora: Ducrotoy (1999) • Plankton: Southward *et al.* (1995); Fromentin & Planque (1996); Nehring (1998); Planque & Taylor (1998); Heath *et al.* (1999); Edwards *et al.* (2001); Beare *et al.* (2002); Beaugrand *et al.* (2002); Edwards *et al.* (2002); Sims & Reid (2002); Beaugrand (2003); Beaugrand & Reid (2003); Edwards & Beaugrand (2007); Greve (2007); Pitois & Fox (2007); Slagstad *et al.* (2007); Voss *et al.* (2007) • Intertidal and benthic organisms: Southward *et al.* (1995); Kroncke *et al.* (1998) • Fish: Scarnecchia (1984); Svendsen (1995); Alheit & Hagen (1997); Southward *et al.* (1988); Sims & Reid (2002); Beaugrand & Reid (2003); Beaugrand *et al.* (2003); Clark *et al.* (2003); Genner *et al.* (2004); Sims *et al.* (2004); Brander (2005); Cotton *et al.* (2005); Perry *et al.* (2005); Rose (2005); Sissener & Bjorndal (2005); Pörtner & Knust (2007); Stenevik & Sundby (2007); Voss *et al.* (2007) • Cephalopods: Sims *et al.* (2001)
Tropical Atlantic	• Plankton: Piontkovski & Castellani (2007)
Antarctic Peninsula	• Plankton: Moline *et al.* (2004)
Bay of Biscay	• Fish: Poulard & Blanchard (2005)
North Pacific Ocean	• Plankton: Roemmich & McGowan (1995); Brodeur *et al.* (1999); Batten (2007)
Worldwide	• Coral reefs: Sebens (1994); Sheppard (1999); Spalding *et al.* (2003)

POTENTIAL EFFECTS OF CLIMATE CHANGE-INDUCED ENVIRONMENTAL VARIATIONS ON MEDITERRANEAN MARINE SPECIES

1. Climate impacts on marine fauna

The global warming-induced alteration of precipitation, temperature, CO_2 concentration and wind patterns will result in a cascade of changes in the physical (e.g. vertical stability of the water column, upwelling regimes, water mass formation and circulation, current patterns), chemical (e.g. seawater pH, salinity, nutrient ratios) and biological (e.g. species phenology, recruitment, physiology, distribution, abundance, diversity, productivity) properties of marine systems (Figure 1) (e.g. Bianchi, 1997; Walther *et al.*, 2002). Temperature anomalies, even over a short period of time, can significantly affect Mediterranean ecosystems and biological diversity (Walther *et al.*, 2002; Anadón *et al.*, 2007). When a keystone species is affected, even by a slight change in climate, the composition and diversity of marine communities can be disrupted (Sanford, 1999). The structure, distribution and phenology of Mediterranean plankton communities, which are at the base of the food chain and which strongly depend on hydroclimatic factors and nutrient ratios, are experiencing change (Velsch, 1997; Turley, 1999; Licandro & Ibanez, 2000; Béthoux *et al.*, 2002; Fernandez de Puelles *et al.*, 2004; Mercado *et al.*, 2005; Molinero *et al.*, 2005a, b; Molinero *et al.*, 2007; Voarino, 2006). For instance, high positive anomalies in water temperature during the 1980s resulted in jellyfish and a drop in abundance of copepods (Molinero *et al.*, 2005a).

Climate has long been recognized as one of the most critical factors influencing Mediterranean resource variability (Garcia & Palomera, 1996). For instance, the most essential environmental mechanisms controlling the growth, abundance, distribution, composition, diversity and recruitment success of Mediterranean species, such as anchovy (*Engraulis encrasicolus*) or sardine (*Sardinella aurita* and *Sardina pilchardus*), include: regional temperature variations, riverine inputs and wind-induced mixing, which influence sea surface temperature and salinity; hydrographical features (e.g. oceanic fronts, water column stability, upwelling zones, sea state); and nutrient enrichment and planktonic production (e.g. Cury & Roy, 1989; Sabates & Gili, 1991; Garcia & Palomera, 1996; Regner, 1996; Agostini & Bakun, 2002; Lafuente *et al.*, 2002; LLoret *et al.*, 2004; Lafuente *et al.*, 2005; Basilone *et al.*, 2006; Sabates *et al.*, 2006).

The Mediterranean Sea is considered oligotrophic because of its low nutrient input from rivers and the nutrient depleted Atlantic water inflow through the Strait of Gibraltar (e.g. Estrada, 1996; Turley, 1999; Zenetos *et al.*, 2002). Winter vertical mixing, coastal upwelling and river runoff are the mechanisms of nutrient input into the euphotic zone, on which marine productivity is dependent. Ocean physical processes such as upwelling phenomena have a major influence on the distribution of primary production, through ascending movements of nutrient-rich deep water into the euphotic zone. Therefore, by altering oceanic features, climate change may affect nutrient availability.

Increasing temperatures are likely to trigger stronger thermal stratification and deepen the thermocline, which could prevent or modify the mixing of water-masses, and cold and nutrient-rich deep waters upwelling (Roemmich & McGowan, 1995). Some areas like the southern Adriatic Sea, where local winter climatic conditions (e.g. winter heat

SC/N10/CC For Info 2



Fig. 1. Main greenhouse gases-related factors influencing marine organisms.

losses and precipitation) strongly influence nutrient availability and plankton production (Gacic, 2002), may be especially vulnerable to climate change.

Extreme meteorological events (e.g. storms and flooding episodes) and sea level rise will release abundant terrestrial suspended solids and pollutants into the marine environment, which may negatively affect coastal biocenoses. Fragile biotopes of the endemic *Posidonia oceanica* have been shown to be vulnerable to physical and chemical damage from meteorological events such as these (Orr *et al.*, 1992; Bombace, 2001). As these sea grass meadows represent a spawning and nursery habitat for numerous species and play a major ecological role, their disappearance would to expected to have significant consequences for coastal ecosystems (Francour, 1997).

By influencing the north-western Mediterranean climate variability (Pozo-Vasquez *et al.*, 2001; Gasparini & Astraldi, 2002; Rixen *et al.*, 2005), the NAO affects the local species composition of planktonic copepods, such as the two dominant copepods of the north-west Meditarranean, *Centropages typicus* and *Temora stylifera* (Molinero *et al.*, 2005a, b).

The high positive NAO episode that caused positive temperature anomalies in the 1980s in the western Mediterranean Sea induced a jellyfish bloom and resulted in a significant diminishment of copepod abundance (Molinero *et al.*, 2005a). This event, which is likely to have been encouraged by global warming, led to high abundance of *Centropages typicus* at the expense of *Temora stylifera* (Molinero *et al.*, 2005b). Jellyfish, which feed on fish larvae, eggs and copepods, can strongly affect plankton communities (Mills, 1995). Changes in planktonic copepods, which affect the fluxes of matter and energy in the marine ecosystem, supply a biological pump of carbon into the deep ocean and strongly influence fish recruitment could alter the ecosystem functioning (Ohman & Hirche, 2001; Molinero *et al.*, 2005b).

Nutrient inputs from the Rhône and Ebre rivers, combined with strong wind mixing, make the north-western Mediterranean highly productive (Lloret *et al.*, 2001). As previously noted, climate change is expected to induce positive NAO episodes with less precipitation and runoff. Consequently, the recruitment of north-western Mediterranean fish, which strongly depend on riverine nutrient supply, may be negatively affected by climate change (Lloret *et al.*, 2001).

In certain areas some organisms survive under specific temperature conditions and cannot adapt themselves or move when the environmental conditions change (e.g. Kenney, 1990; MacGarvin & Simmonds, 1996; Hughes, 2000). Dispersal limitation can limit the response of marine animals such as benthic organisms that could sometimes be unable to successfully migrate toward more suitable environments (problems would include long distances or travelling against strong currents) (Hiscock *et al.*, 2004).

By affecting the physiology of marine organisms, global warming could impact the performance and survival of those organisms that live close to their thermal tolerance or situated at the northern or southern limit of their distribution (Laubier, 2001; Hochachka & Somero, 2002; Somero, 2002; Poulard & Blanchard, 2005). Similarly, some larval and young benthic stages of some organisms are more sensitive to temperature than adults (Foster, 1971; Pechenik, 1989). According to Bella Galil in Cheviron (2007), deep-water organisms that live in constant temperatures (~13°C) and that are not used to seasonal temperature variation will be vulnerable to climate change.

SC/N10/CC For Info 2

2. Climate impacts on marine flora

Climate change is likely to affect phytoplankton composition by affecting nutrient concentrations and ratios. Over the last two decades, in the north-western Mediterranean, meteorological anomalies (e.g. warmer water, decreased salinity, longer periods of sunshine and lower wind stress) have affected water column stability and reduced nutrient replenishment into the euphotic zone (Goffart *et al.*, 2002). This event decreased silicon availability, which in turn triggered a reduction of diatom abundance and a shift toward non-siliceous species, such as flagellates and dinoflagellates (Turley, 1999; Béthoux *et al.*, 2002; Goffart *et al.*, 2002). By modifying the composition of phytoplankton communities, climate change could then seriously alter nutrient cycling and food web dynamics (Litchman *et al.*, 2006).

Warmer temperatures trigger changes in the timing of plankton blooms, resulting in a temporal mismatch between primary production and higher trophic levels of the food web (Edwards & Richardson, 2004; Hiscock *et al.*, 2004). In north-western European estuaries, for example, a long-term data set (1973–2001), showed an increase of spring temperatures of 0.07°C yr^{-1}, which resulted in the earlier spawning of a bivalve, *Macoma balthica*, but did not affect phytoplankton blooms, causing a divergence in timing between larval bivalve production and prey (Philippart *et al.*, 2003). In addition, rising seawater temperatures advance the onset of crustacean reproduction and enhance shrimp predation pressure on vulnerable juvenile spat that leads to low recruitment success of the spat (Philippart *et al.*, 2003).

MIGRATION OF SPECIES AND INCREASED NUMBER OF EXOTIC THERMOPHILIC SPECIES IN THE MEDITERRANEAN SEA

The presence of exotic species in the Mediterranean has resulted from the combination of environmental factors and human activities like the opening of the Suez Canal in 1869. Today, more than 500 alien species are recorded in the Mediterranean Sea and their geographical ranges are increasing, as is the rate of increase of new alien species being identified (Galil & Zenetos, 2002; Zenetos *et al.*, 2003; Harmelin-Vivien *et al.*, 2005; CIESM, 2005; Galil, 2007). Climate change, combined with the establishment of exotic species has led to the 'tropicalization' of the Mediterranean (Bianchi, 2007). The increase of alien species can cause endemic species to rapidly decline in abundance and be displaced (Galil & Zenetos, 2002; Zenetos *et al.*, 2002; Galil, 2007). Such phenomena can alter the infra-littoral communities and induce ecological impacts such as local population decline and extirpation, reduction of genetic diversity in native species, foodweb alterations, loss of habitat functions, processes and structure, increase in the risk of extinction and biotic homogenization (Ricciardi, 2004; Galil, 2007).

In the North Atlantic Ocean, the observed northward migration of species (250 km per decade), resulting from a minor increase in temperature, is likely to be linked to global warming (Parmesan & Yohe, 2003; EEA, 2004; Oviatt, 2004). Perry *et al.* (2005) have also demonstrated that, in the North Sea, recent increases in sea temperature have led to nearly two-thirds of fish species (exploited and non-exploited) shifting in mean latitude or depth or both over 25 years. For species with northerly or southerly range margins in the North Sea, half showed boundary shifts with warming, and all but one species shifted northward.

In the Mediterranean, over the past three decades, increasing water temperature has been observed in the Ligurian Sea (Béthoux *et al.*, 1990; Astraldi *et al.*, 1995), which is one of the coldest zones of the Mediterranean. This phenomenon encouraged warm-water species to shift their ranges northward and settle in the Ligurian waters where they were formerly rare or absent (Bianchi & Morri, 1993, 1994; Francour *et al.*, 1994; Astraldi *et al.*, 1995; Morri & Bianchi, 2001).

Consequently, warm-water species like the ornate wrasse (*Thalassoma pavo*) colonized and established large and stable populations in the north-western Mediterranean (Bianchi & Morri, 1994; Bombace, 2001; Vacchi *et al.*, 2001). Other thermophilic species like the grey triggerfish (*Balistes carolinensis*), Mediterranean parrotfish (*Sparisoma cretense*), round sardine (*Sardinella aurita*), bluefish (*Pomatomus saltatrix*), Senegalese sole (*Solea senegalensis*), dusky grouper (*Epinephelus marginatus*), bastard grunt (*Pomadasys incisus*), European barracuda (*Sphyraena sphyraena*), slackskin blaasop (*Sphoeroides cutaneus*), the coral *Astroides calycularis* and groupers of the genus *Epinephelus* have been frequently recorded amongst the 'cold biota' of the northern Mediterranean (Vacchi & Cau, 1986; Serena & Silvestri, 1996; Relini & Orsi Relini, 1997; Dulcic *et al.*, 1999; Louisy & Culioli, 1999; Dulcic & Grbec, 2000; Guidetti & Boero, 2001; Dulcic *et al.*, 2005, 2006; Athanassios & Antonopoulou, 2006).

The recent northward expansion of tropical groupers, like the white grouper (*Epinephelus aeneus*) and dusky grouper (*Epinephelus marginatus*), may result from the warming of Mediterranean waters (Francour *et al.*, 1994; Dulcic & Lipej, 1997; Zabala *et al.*, 1997). Since groupers are top carnivores and among the bigger coastal fish species, their successful colonization is likely to affect the ecology of endemic species and influence local fisheries (Glamuzina, 1999).

Similarly, in the Adriatic Sea, the presence of thermophilic species of fish and zooplankton, that were formerly uncommon or absent in this zone, has increased in the past 30 years (Dulcic & Grbec, 2000; Kamburska & Fonda-Umani, 2006). These observations have been correlated with seawater warming and salinity changes occurring since 1988 (Francour *et al.*, 1994; Russo *et al.*, 2002) and, thus, global warming is thought to be responsible for the observed faunal changes in this region (Dulcic *et al.*, 1999; Dulcic & Grbec, 2000; Parenti & Bressi, 2001).

The northward migration of warm water species will induce species competition for existing niches (IPCC, 2001a) and thermophilic species are likely to increase in abundance at the expense of cold-water species (Beaugrand *et al.*, 2002; Galil & Zenetos, 2002; Hiscock *et al.*, 2004). Marine bioinvasions, which are considered to be 'biological pollution' (Bouderesque & Verlaque, 2002; Elliot, 2003) have altered marine ecosystems (e.g. competition with indigenous species, food web shifts) and are considered as one of the most intense and damaging anthropogenic impacts (Harris & Tyrrell, 2001; Frank *et al.*, 2005; Galil *et al.*, 2007). The Mediterranean Sea is one of the most impacted seas of the world in terms of biological invasion (Galil, 2007) and climate change is likely to facilitate invasion of thermophilic alien species causing irreversible impacts on native populations (Carlton, 2000; Stachowicz *et al.*, 2002b; Gritti *et al.*, 2006; Galil *et al.*, 2007; Occhipinti-Ambrogi, 2007). According to Bella Galil (in Cheviron, 2007) alien species like the bivalve *Brachidontes pharaonis* or the jellyfish *Rhopilema nomadica*, both belonging to the 100 'worst invasives' species, could

SC/N10/CC For Info 2

lead to the extinction of numerous native Mediterranean species. Photophilic subtidal macrophyte assemblages appear particularly vulnerable to invasions of exotic algal species such as *Caulerpa taxifolia* (Streftaris & Zenetos, 2006).

In addition to northward migration, bathymetric displacements occur among populations of invasive and endemic species (Galil & Zenetos, 2002). This is the case for the indigenous red mullet (*Mullus barbatus*) and hake (*Merluccius merluccius*) with both moving into deeper and cooler waters due to their respective warm-water competitors: the goldband goatfish (*Upeneus moluccensis*) and brushtooth lizardfish (*Saurida undosquamis*) (Oren, 1957). Similarly, in the north-western Mediterranean, the endemic spottail mantis shrimp (*Squilla mantis*) is usually observed in deeper waters (70–80 m) rather than the thermophilic Red Sea mantis shrimp (*Oratosquilla massavensis*) (10–25 m) (Galil & Zenetos, 2002).

The warming of the Mediterranean waters may modify species' migration periods causing changes in the trophic webs. For example, Bombace (2001), has documented that in the last few decades, the amberjack (*Seriola dumerilii*) and bluefin tuna (*Thunnnus thynnus*) appear to stay longer (until mid-winter instead of autumn) in the northern and central Mediterranean before migrating toward their winter territories.

Although there are presently no marine species extinctions that have been correlated with global warming, today, species such as the Mediterranean mysid *Hemimysis speluncola* are regarded as threatened (Chevaldonné & Lejeusne, 2003; Lejeusne, 2005; Lejeusne & Chevaldonné, 2005; Harley *et al.*, 2006). Furthermore, Chevaldonné & Lejeusne (2003) showed that the endemic cave-dwelling invertebrate *Hemimysis speluncola* which was previously abundant in the north-western Mediterranean, has been replaced by a warm-water species (*Hemimysis margalefi*).

Areas of high endemic biodiversity are likely to be less subject to non-indigenous species invasion (Kennedy *et al.*, 2002; Stachowicz *et al.*, 2002a; Duffy, 2003). Therefore, sparse and declining populations such as *Posidonia* meadows are more vulnerable to be overgrown and replaced by invasive macroalgae like *Caulerpa taxifolia* and *Caulerpa racemosa* (Meinesz *et al.*, 2001; Bouderesque & Verlaque, 2002; Peirano *et al.*, 2005). Climate change could be one of the factors responsible for the decline of *Posidonia* and for the expansion of *Caulerpa* (Komatsu *et al.*, 1997; Raniello *et al.*, 2004; Peirano *et al.*, 2005; Ruitton *et al.*, 2005).

Furthermore, some cold-water species like the small euphausid species *Meganyctiphanes norvegica*, which is situated at the northern limit of its ecological tolerance, would be more vulnerable to invasion. This is especially true for many taxa in the eastern Mediterranean, leaving this region more exposed to invasion (Galil & Zenetos, 2002).

LARGE MASS-MORTALITY EVENTS AND DECLINE IN SPECIES ABUNDANCE AND DIVERSITY ASSOCIATED WITH TEMPERATURE ANOMALIES

Temperature anomalies have led to several mass-mortality episodes in the Mediterranean Sea. In 1999, successive heat waves with consequent peaks in water temperature and a deepening of the thermocline caused a mass-mortality event of 28 invertebrate species in the north-western Mediterranean (Cerrano *et al.*, 2000; Perez *et al.*, 2000; Romano *et al.*, 2000; Laubier *et al.*, 2003). This event affected benthic organisms such as gorgonians, sponges, cnidarians, bivalves, ascidians, bryozoans, scleractinian corals and zoanthids (Cerrano *et al.*, 2000; Perez *et al.*, 2000; Romano *et al.*, 2000; Garrabou *et al.*, 2001). According to Occhipinti-Ambrogi (2007), niches resulting from mass mortality events become available for new invasive colonizers.

Variations of the nutrient load and seawater properties (e.g. temperature increase) can lead to coastal eutrophication and algal bloom (Degobbis *et al.*, 2000). This phenomenon particularly concerns the shallow northern Adriatic waters (UNEP, 1996; Degobbis *et al.*, 2000). By increasing seawater stratification, meteorological anomalies regularly cause bottom water anoxia and red tide events in this region, provoking mass mortality episodes of fish and benthic organisms (Degobbis *et al.*, 2000; Anadón *et al.*, 2007). Eutrophication events occurred frequently during the second half of the 20th Century (Degobbis *et al.*, 2000) and the intensity, frequency and geographical expansion of algal blooms became a growing concern since the 1970s in this area (Justic, 1987). According to Boero (1996), increases in jellyfish (*Pelagia noctiluca* and *Aurelia aurita*), salps (Thaliacea), harmful algal blooms and red tides were all promoted by abnormal meteorological and oceanographic changes occurring since 1988 in the Adriatic Sea.

During the last decades, the collapse of sprat (*Sprattus sprattus*) and anchovy (*Engraulis encrasicolus*) stocks in the Adriatic affected the species stocks of the entire Mediterranean and was associated with the decrease of surface temperature resulting from climatic anomalies (Regner, 1996; Salat, 1996; Bombace, 2001; Azzali *et al.*, 2002). Similarly, during the 2001 winter, climate-induced low surface temperatures led to the decline of sardines (*Sardinella aurita*) and phytoplankton blooms (Guidetti *et al.*, 2002).

By altering oceanographic properties like seawater temperature, salinity, water transparency and deep water oxygen saturation, climate change could impact the entire Adriatic Sea ecosystem (Zore-Armanda *et al.*, 1987; Zore-Armanda, 1991; Russo *et al.*, 2002).

Global warming may promote the development of toxic dinoflagellates like *Gymnodinium catenatum* which is responsible for frequent Mediterranean toxic events, causes paralytic shellfish poisoning (PSP) and may alter the marine ecosystem (Garcés *et al.*, 2000; Taleb *et al.*, 2001; Vila *et al.*, 2001; Calbet *et al.*, 2002; Gomez, 2003). According to Calbet *et al.* (2002), the occurrence of *Gymnodinium catenatum* (toxic dinoflagellate) in the Alboran Sea reduced the grazing impact of mesozooplankton on the microbial communities and may have altered the Mediterranean pelagic food web.

Gomez & Claustre (2003) and Polat (2004) suggest that the presence of new warm-water dinoflagellate species in the Mediterranean Sea, like *Asterodinium libanum*, *Asterodinium gracile* and *Citharistes regius* is likely to be associated with the warming of Mediterranean waters. A species of a similar genus, *Ostreopsis armata*, was recently observed (summer 2005 and 2006) in the Ligurian Sea where it repeatedly bloomed and apparently triggered respiratory diseases in humans (details in Occhipinti-Ambrogi, 2007).

Meteorological anomalies can significantly alter ecosystems and cause mass mortality episodes. In the eastern Mediterranean Sea, a climatic event, called the Eastern Mediterranean Transient (EMT), was correlated with local meteorological anomalies (reduced precipitation, change in wind patterns and cold winters) (Klein *et al.*, 1999) and resulted in a drastic alteration of faunal abundance and diversity (Danovaro *et al.*, 2001). By modifying the physico-chemical

SC/N10/CC For Info 2

characteristics of the deep waters, this temperature shift significantly and rapidly affected deep-sea nematode diversity (Danovaro *et al.*, 2004). Following the 1994–1995 period, when the temperature recovered, only some marine faunal species recovered (Danovaro *et al.*, 2004). These observations give us a better vision of the potential large-scale consequences of global warming.

INCREASE OF DISEASES AND PATHOGENS

Since warmer temperatures are known to favour the presence of pathogens, epidemiological outbreaks are likely to become more severe and frequent with the warming of Mediterranean waters (Gantzer *et al.*, 1998; Harvell *et al.*, 1999, 2002; Marcogliese, 2001; Drake *et al.*, 2007). According to CIESM (2004) global warming could lead to the development of tropical and subtropical pathogens across the Mediterranean Sea. Along the Ligurian Sea, the massive development of a cyanobacterium combined with warm water, caused several populations of the zoanthid *Parazoanthus axinellae* to decline since 2000 (Cerrano *et al.*, 2006). This species has been replaced by an incrusting thermophilic sponge (*Crambe crambe*) which quickly colonized the niche deserted by *Parazoanthus* (Cerrano *et al.*, 2006).

Similarly, the thermophilic bacteria *Vibrio shiloi*, was involved in the mortality episode of the Mediterranean coral *Oculina patagonica* (Kushmaro *et al.*, 1998) and the 1999 mass mortality event in the Ligurian Sea was induced by the combination of a temperature shift with the growth of opportunistic warm-water pathogens (Cerrano *et al.*, 2000).

Simmonds & Mayer (1997) provided a tentative link between reduced nutrient input to the western Mediterranean basin (resulting from reduced rain fall) and the initiation of the striped dolphin (*Stenella coeruleoalba*) mass mortality in 1990. This is discussed further below.

OTHER EXPECTED CHANGES

Increased concentration of atmospheric anthropogenic CO_2 and climate change is likely to alter marine ecosystems in several other ways. As temperature increases, the oxygen solubility decreases and fish metabolism accelerates (Green & Carritt, 1967; Pörtner & Knust, 2007). Since the demand for oxygen and food will be enhanced in order to support higher metabolic rates, the decreased concentration of dissolved oxygen will thus affect fish growth and breeding capacity, and could lead to the extinction or migration (to cooler waters) of some fish species (Pörtner & Knust, 2007). Furthermore, enhanced fish metabolic rates and food consumption resulting from higher seawater temperatures may increase fish pollutant uptake (e.g. mercury) which will then be transferred into higher levels of the food chain (Harris & Bodaly, 1998).

In addition, the rise of atmospheric CO_2 could increase the acidity of seawater and therefore reduce the saturation state of $CaCO_3$ species in the oceans, namely calcite and aragonite (Caldeira & Wickett, 2003; Orr *et al.*, 2005). The marine species most likely to be affected by this acidification will be small and thin-shelled organisms that use $CaCO_3$ such as calcifying plankton (e.g. coccolithophores), coralline algae, pteropod molluscs and coral polyps (e.g. reef-building scleractinian corals) (Kleypas *et al.*, 1999; Riebesell *et al.*, 2000; Feely *et al.*, 2004). By reducing the level of calcium carbonate saturation, ocean acidification will affect the process of calcification of some marine key organisms. Feely *et al.* (2004) argue that the calcification rate of multiple taxa will be affected, from single celled protists to reef-building corals—and across all CaCO mineral phases. Cephalopods may also be particularly sensitive and this is described in Table 2 at the end of this paper. According to a study recently carried out by The Royal Society (2005), the plankton calcification process could become very limited and a large portion of marine life could then disappear by 2100.

The seawater acidification phenomenon, which is responsible for nanism (a genetic anomaly resulting in short stature) and malformation symptoms in several phytoplankton species (Feely *et al.*, 2004), can also have an impact on the reproductive patterns of fish (Stanley, 1984).

According to Caldeira & Wickett (2003) and Feely *et al.* (2004), the expected change in pH will be greater than any other pH variations observed in the fossil record over the last 200–300 million years. Manipulative experiments showed that a three-month reduction of pH by 0.7-unit reduced mussel metabolism and growth (Michaelidis *et al.*, 2005). Similarly, a six-month, 0.03-unit pH reduction, which corresponds to a 200-ppm increase in atmospheric CO_2, lowered gastropod and sea urchin growth and survival (Shirayama & Thornton, 2005). A study by Alvarez *et al.* (2005) shows that, although ocean uptake of carbon dioxide for the Mediterranean appears small in terms of global ocean uptake, 'the impact on local carbonate chemistry will be large.'

In addition to this process of ocean acidification, changes in marine ecosystems due to global warming (e.g. water temperature, algal bloom enhancement and water colour) might affect the visual sensitivity of fish (spectral sensitivity) (Archer *et al.*, 2001).

Several points discussed in this section are likely to be relevant to a broad variety of marine species including key species in Mediterranean ecosystems, with significant consequences for foodweb structures, marine system functions and ecosystem equilibrium (Petchey *et al.*, 1999; Sanford, 1999; Schiel *et al.*, 2004). Such foodweb alterations are likely to produce significant cascade effects on marine biodiversity including impacts on species of higher trophic levels, such as cetaceans.

Climate change impacts on cetaceans

The distribution of cetaceans, which have a major influence on marine community function and structure (e.g. Katona & Whitehead, 1988; Bowen, 1997; Jones *et al.*, 1998), is closely related to environmental parameters such as oceanographic features and food availability (Millot & Taupier-Letage, 2004). According to MacGarvin & Simmonds (1996), they are not likely to be able to adapt to rapid shifts in temperatures and environmental conditions, and climate change may represent the most serious long-term threat to cetaceans (Burns, 2001). Although lower trophic levels are the most likely to be altered, cetaceans could be affected by global warming in a variety of ways (Figure 2; Table 2) (e.g. Reeves, 1991; Fisher *et al.*, 1994; Agardy, 1996; IWC, 1997; European Community, 1999; Hardwood, 2001; Simmonds & Nunny, 2002; Gambaiani *et al.*, 2005; Learmonth *et al.*, 2006; Simmonds & Isaac, 2007).

CHANGE IN FOOD SUPPLY

The distribution, abundance and migration of cetaceans is strongly influenced by prey availability (e.g. Kenney *et al.*, 1996) and cetaceans which are confined in restricted habitat,



Fig. 2. Overview of the potential impacts of global warming on marine life, including cetaceans (from Gambaiani *et al.*, 2005).

with limited ranges, are likely to be most vulnerable to climate change (e.g. Learmonth *et al.*, 2006; Simmonds & Isaac, 2007).

Change in key prey species distribution is the main driving factor defining geographical range and habitat preference in cetaceans (e.g. Evans, 1971; Wells *et al.*, 1990; Hanson & Defran, 1993; Simmonds, 1994; Agardy, 1996; Maze & Würsig, 1999).

For instance, in the eastern North Pacific Ocean, an increase in seawater temperature combined with a change in oceanographic conditions is thought to have led to the death of hundreds of grey whales (*Eschrichtius robustus*) as the result of a decline in their prey species (Grebmeier & Dunton, 2000; Moore *et al.*, 2003; Gulland *et al.*, 2005). In the Mediterranean Sea, the decline of several cetacean populations has been associated with the reduction of prey resources (Perrin, 1989; UNEP/IUCN, 1994; Reeves *et al.*, 2003; Reeves & Notarbartolo di Sciara, 2006).

Odontocete prey species such as pilchard (*Sardina pilchardus*) are affected by climate change (Southward *et al.*, 1988; Garcia & Palomera, 1996; Regner, 1996; Bearzi *et al.*, 2003). Pilchards have been shown to be a key prey species for common dolphins off the Portuguese coast (Silva, 1999). In addition, cephalopods, which represent the main food supply for numerous Mediterranean cetacean species (Wurtz & Marrale, 1991; Bompar, 2000; Bearzi *et al.*, 2003) seem to be particularly vulnerable to environmental changes including pH and temperature (Sims *et al.*, 2001; Pierce & Boyle 2003; Arkhipkin *et al.*, 2004) (Table 2).

In the Adriatic Sea, climatic shifts are suggested to have altered the distribution of the key prey species of common dolphins (*Delphinus delphis*) and bottlenose dolphins (*Tursiops truncatus*) (Blanco *et al.*, 2001; Bearzi *et al.*, 2003). In particular, the climate-induced increase in abundance of thermophilic species such as round sardinella and jellyfish may have caused the European anchovy (*Engraulis encrasicolus*) population to decrease (Regner, 1996).

Shifts in prey species availability may force cetaceans to change their feeding strategies and spend more time and energy foraging, which could have drastic consequences on their health and could affect their immune systems (Northridge, 1984; Shane, 1990; Bräger, 1993; Smith & Whitehead, 1993; Agardy, 1996; Stern, 1996; Bearzi, 2002). A high proportion of time and effort devoted to feeding-related activities was recorded in Mediterranean bottlenose dolphins in the northern Adriatic Sea as a response to environmental changes and reduced prey availability (Politi, 1998; Bearzi *et al.*, 1999). Consequently, the time dedicated to socializing and breeding is reduced, with negative consequences on cetacean reproductive success (Valiela, 1995; Bearzi, 2002). The fact that climate change impacts can lead to reduced prey availability and subsequently affect the health, physical strength and abundance of cetacean populations, has been observed in bottlenose dolphins in the eastern Ionian Sea (Politi *et al.*, 2000; Politi & Bearzi, 2004).

Aguilar & Raga (1993) and Simmonds & Mayer (1997) have suggested that the mass mortality of thousands of striped dolphins (*Stenella coeruleoalba*) during the 1990–1992 morbillivirus epizootic might have been caused by the unusual warm and dry winter of 1989–1990 that led to abnormally warm water temperatures and low rainfall. This phenomenon resulted in reduced nutrient input into the eastern Mediterranean and thus low productivity (Simmonds & Mayer, 1997). This led to the decline of the dolphin's common prey and explains why many of the dolphin carcasses showed depleted body fat reserves (Aguilar *et al.*, 1991).

Similarly, during a survey carried out in Corsica, Dhermain (2003) observed fewer bottlenose dolphins in coastal waters than usual. The excessively hot 2003 summer, which resulted in abnormally high coastal water temperatures, could explain the migration of bottlenose dolphins toward the open sea. Such exceptional meteorological events may illustrate how an increase of temperature could impact on cetaceans.

Moreover, global warming is likely to encourage the spreading of viruses and pathogens and may promote epizootic events like morbillivirus infections (Agardy, 1996), which have also been identified in the endangered

SC/N10/CC For Info 2

Table 2. Impact of climate changes on cetaceans around the world.

Global warming-induced changes	Impacts on marine mammals with some examples of species most likely to be, or, already affected	
Seawater temperature	Directly affected: • Endangered, young animals, species with low mobility, restricted distribution, low thermal tolerance like vaquitas (*Phocoena sinus*) in the Gulf of Mexico; Arctic bowhead whales (*Balaena mysticetus*); Arctic belugas (*Delphinapterus leucas*); narwhals (*Monodon monoceros*); finless porpoises (*Neophocaena phocaenoides*), tropical dolphins, humpback whales (*Megaptera novaeangliae*) in the Indian Ocean, Mediterranean fin whales (*Balaenoptera physalus*) and river dolphins (IWC, 1997; Bannister, 2002; Würsig *et al.*, 2002; Simmonds, 2004; COSEWIC, 2005; Laidre & Heide-Jorgensen, 2005; Robinson *et al.*, 2005; Learmonth *et al.*, 2006; Simmonds & Isaac, 2007) • Animals like bottlenose dolphins (*Tursiops truncatus*) in the Gulf of Mexico, living in coastal zones where seawater temperature variation is enhanced (IWC, 1997) • Cold-water species like white-beaked dolphins (*Lagenorhynchus albirostris*) in north-western Scotland whose relative abundance and occurrence decreased whereas warm-water common dolphins (*Delphinus delphis*) increased in this region (MacLeod *et al.*, 2005) • Long-finned pilot whale (*Globicephala melas*) population structure in the North Atlantic and short-finned pilot whales (*Globicephala macrorhynchus*) distribution in Japan (Fullard *et al.*, 2000) Effects on prey availability affecting the distribution and geographical range of: • Long-finned pilot whales in the Faeroe Islands (Bjorge, 2002) • Bottlenose dolphins in north-east Scotland (Wilson *et al.*, 2004) Effects on prey availability affecting the reproductive success of: • North Atlantic right whales (*Eubalaena glacialis*) (Greene & Pershing, 2004)	• North-east Atlantic fin whales (Lockyer, 1986) • Humpback whales (Wiley & Clapham, 1993) • Harbour porpoises (*Phocoena phocoena*) in the Bay of Fundy (Read & Gaskin, 1990) Effects on reduced prey availability leading to: • Increased vulnerability to disease particularly for species at the limit of their thermal tolerance (Würsig *et al.*, 2002; Lafferty *et al.*, 2004) • Increase of mass mortality events like morbillivirus that affected Mediterranean striped dolphins (*Stenella coeruleoalba*) (Aguilar & Raga, 1993; Cebrian, 1995; Harvell *et al.*, 1999; Kennedy, 1999; Geraci & Lounsbury, 2002) • Higher exposure to contaminants stocked in blubber and mobilized during starvation (Aguilar *et al.*, 1999), that alter reproductive, endocrine and immune systems (Fuller & Hobson, 1986; Aguilar & Borrell, 1994; Ross *et al.*, 2000) • Stranding of hundreds of grey whales (*Eschrichtius robustus*) in eastern America (Grebmeier & Dunton, 2000; Moore *et al.*, 2003; Gulland *et al.*, 2005) • Episodic shifts in cetacean populations' abundance and distribution in the Gulf of Maine. Concerned species: humpback and fin whales replaced by right and sei whales (*Balaenoptera borealis*); white-beaked dolphins (*Lagenorhynchus albirostris*) replaced by Atlantic white-sided dolphins (*Lagenorhynchus acutus*); and harbour porpoise (*Phocoena phocoena*) (Kenney *et al.*, 1996; Palka *et al.*, 1997) • Episodic shifts in the inshore incidence of Pacific white-sided dolphins (*Lagenorhynchus obliquidens*) in British Columbia (Morton, 2000) • Disappearance of common dolphins (*Delphinus delphis*) along the north-eastern coast of Florida (Caldwell & Caldwell, 1978)
Precipitation and extreme weather occurrence	Effects of warmer seawater temperature and nutrient enrichment due to increased runoff: • Increase in the frequency and strength of toxic algal bloom episodes triggering lethal poisoning in marine mammals (Hernández *et al.*, 1998; Burns, 2002; Geraci & Lounsbury, 2002) • Increase in eutrophication events affecting marine organisms dynamics like phytoplankton in the North Sea (Edwards *et al.*, 2001)	Effects of pollutants inputs into coastal waters due to increased runoff and flooding events: • Impacts on coastal marine mammals and other organisms (including prey species) (e.g. MacGarvin & Simmonds, 1996; Orr *et al.*, 1992; Learmonth *et al.*, 2006) Effects on oceanic properties (coastal waters salinity, circulation, etc) affecting: • The distribution and abundance of cetaceans' prey species (e.g. MacGarvin & Simmonds, 1996; Learmonth *et al.*, 2006)
Sea level rise	Flooding of coastal habitats and impacts on species that depend on coastal areas (as breeding, nursing, feeding, mating, resting zones) like: • Mediterranean monk seals (*Monachus monachus*) breeding in caves or on small beaches (Harwood, 2001; Würsig *et al.*, 2002)	• Grey and humpback whales (IWC, 1997)
Ocean circulation	Changes in current patterns and oceanic fronts directly affecting: • The distribution of the majority of tropical and temperate cetaceans (Worms *et al.*, 2005; Learmonth *et al.*, 2006) like sperm whales associated with the Antarctic convergence in the Southern Ocean (Boyd, 2002)	Changes in water mass formation, mixing and upwelling pattern affecting: • Marine biodiversity and ecosystems (IPCC, 2001a) • The distribution, abundance and migration of plankton, fish and cephalopods (e.g. Planque & Taylor, 1998; Waluda *et al.*, 2001; Walther *et al.*, 2002) affecting cetaceans like minke whales (Bjorge, 2002)

Continued

SC/N10/CC For Info 2

Table 2. *Continued.*

Global warming-induced changes	Impacts on marine mammals with some examples of species most likely to be, or, already affected	
Salinity	Direct affects on cetaceans epidermis generating: • More important skin lesions in bottlenose dolphins living in salted and cold waters (Learmonth *et al.*, 2006) • Animal increased stress and susceptibility to diseases or anthropogenic impacts (Wilson *et al.*, 1999)	Effects on prey species through changes in oceanic properties or low salinity tolerance like: • Most cephalopod species (e.g. De Heij & Baayen, 2005) • Coastal phytoplankton of the Antarctic peninsula (Moline *et al.*, 2004)
Seawater pH	Increased CO_2 affecting metabolic function, growth and reproduction of water-breathing organisms (e.g. Pörtner *et al.*, 2004; Orr *et al.*, 2005; Royal Society, 2005)	• High metabolic rate species like ommastrephid squids (e.g. *Illex illecebrosus*) (Learmonth *et al.*, 2006)
Large-scale atmospheric patterns occurrence and strength (e.g. North Atlantic Oscillation (NAO), El Niño, Southern Oscillation, Pacific Decadal Oscillation (PDO))	Increase in frequency of large-scale atmospheric events (Timmermann *et al.*, 1999) affecting the distribution, growth, abundance and recruitment of marine organisms (IPCC, 2001a; Stenseth *et al.*, 2002) including: • Market squid (*Loligo opalescens*), a short-finned pilot whales' prey (Shane, 1995) • North Atlantic cod (*Gadus morhua*) (Stenseth *et al.*, 2002) • North Atlantic copepods *Calanus finmarchicus* and *Calanus helgolandicus* (Planque & Taylor, 1998; Beare *et al.*, 2002; Beaugrand & Ibanez, 2004) Large-scale atmospheric events' impacts on prey availability and thus affecting: • Community structure of short-finned pilot whale (*Globicephala macrorhynchus*) being replaced by Risso's dolphins (*Grampus griseus*) in southern California (Shane, 1995; Würsig *et al.*, 2002) and in the Gulf of Mexico (Jefferson & Schiro, 1997)	• Sperm whale (*Physeter macrocephalus*) reproductive success in the Galapagos (Whitehead, 1997) • Dusky dolphins (*Lagenorhynchus obscurus*) reproductive success in Peru (Manzanilla, 1989) • Short-beaked common dolphins (*Delphinus delphis*) and other Delphinidea distribution in New Zealand (Gaskin, 1968; Neumann, 2001) • Southern right whales breeding success (Leaper *et al.*, 2006) • Californian coastal bottlenose dolphins geographical range (Wells *et al.*, 1990; Würsig *et al.*, 2002) • Grey whale high mortality rate and low recruitment in the Pacific Ocean (Le Boeuf *et al.*, 2000) • Pacific killer whales (*Orcinus orca*) and Atlantic bottlenose dolphins social organization and behaviour (Lusseau *et al.*, 2004) • North Atlantic right whales breeding success (Simmonds & Isaac, 2007) • Sperm whale in the north-east Atlantic (Robinson *et al.*, 2005) • Probably, the North Atlantic cetaceans in general, in response to the NAO (Learmonth *et al.*, 2006)

Mediterranean monk seal (*Monachus monachus*) (Van de Bildt *et al.*, 1999). A worldwide increase in mass mortality events in marine mammals has been reported by Simmonds & Mayer (1997).

It is interesting to note that, as cetaceans are long-lived, slow-reproducing animals (generally producing one offspring per female every 2–3 years), when a population is severely diminished by a virus or other agents, recovery may be slow and such species can relatively easily become endangered (Dhermain *et al.*, 2002; Reeves *et al.*, 2003).

According to Greene & Pershing (2004), in the North Atlantic Ocean, the effects of global warming on the abundance of *C. finmarchicus* strongly influence right whale (*Eubalaena glacialis*) calving rates. A similar situation is likely to take place in the Mediterranean. For instance, *Meganyctiphanes norvegica*, which constitutes the only known food supply of the fin whales (*Balaenoptera physalus*) in this region, is at the northern limit of its distribution (Besson *et al.*, 1982; Viale, 1985; Orsi Relini & Giordano, 1992; Orsi Relini *et al.*, 1994; Gannier, 1995, 1997; Forcada *et al.*, 1996; Astruc & Beaubrun, 2001; Notarbartolo di Sciara *et al.*, 2003) and thus, in case of unsuitable environmental properties, will not be able to move northward because of the physical land barrier.

The distribution and abundance of this euphausid are correlated with specific hydrobiological parameters (e.g. seawater temperature, salinity, food availability and current patterns) (Pustelnik, 1976; De la Bigne, 1985; Macquart-Moulin & Patriti, 1996; Velsch, 1997). Climate change-induced alteration of ocean circulation is likely to modify larval transport processes, species dispersal and recruitment, and impact krill population dynamics (Harley *et al.*, 2006).

According to Einarsson (1945), in the north Atlantic, the optimal temperature range of *Meganyctiphanes norvegica* is between 2 and 15°C with a high mortality rate above 15°C (Buchholz *et al.*, 1995). This thermal limit is 18°C for the Mediterranean population (Fowler *et al.*, 1971). Salinity change is also likely to affect this species which has its tolerance limit at 20–24 ppm (Forward & Fyhn, 1983).

Whereas, temperature warming in the north-east Atlantic has led to the migration of several marine organisms to northern latitudes (e.g. Beaugrand *et al.*, 2002), in the Mediterranean Sea, *Meganyctiphanes norvegica* will not be able to extend its range northward because of the land barrier and is likely to share its environment with more thermophilic invasive species in the future.

Furthermore, calcifying organisms including some phyto- and zooplankton species are likely to be affected by acidification (Royal Society, 2005) and a possible temporal mismatch may result between *Meganyctyphanes norvegica* and phytoplankton blooms, its food supply, with protential consequences for predators including the endangered bluefin tuna (*Thunnus thynnus*), albacore tuna (*Thunnus alalunga*) (Quynh, 1978), squid (*Illex coindetii*) (Sanchez, 1982) and the Mediterranean fin whale population.

Mediterranean fin whale distribution can be expected to be affected by food availability (Littaye *et al.*, 2004) and since

SC/N10/CC For Info 2

Mediterranean fin whales are genetically and reproductively isolated from those of the Atlantic (Bérubé *et al.*, 1998), they are regarded as more vulnerable to environmental pressures, including global warming (Dhermain *et al.*, 2002).

By affecting the distribution and abundance of cetacean prey species, climate change is likely to trigger dietary competition between species, and could then cause inter- and intra-species competition between Mediterranean cetaceans. This is particularly true for striped dolphins and common dolphins (Aguilar, 2000), whose food web dynamics have been affected by recent temperature changes (Bearzi *et al.*, 2003).

In addition, a decrease of prey species may increase cetacean mortality rates and vulnerability to diseases as a consequence of reduced immune function, as in the Mediterranean striped dolphin epizootic outbreak in the 1990s (Aguilar & Raga, 1993; Bearzi, 2002).

CLIMATE CHANGE COMBINED WITH FISHERIES PRESSURE

The combination of climate-induced impacts with other anthropogenic impacts like overfishing is likely to impact cetaceans (CIESM, 2000; Bearzi, 2002). Several cetacean species such as coastal bottlenose dolphins and common dolphins are already competing with fishermen for prey species exploited by fisheries (Bearzi, 2002; Abad *et al.*, in press). As previously observed prey species distribution and abundance could be severely affected by global warming. The diminution of fish stocks is likely to result in a stronger competition with fishermen and in higher risks of harassment of dolphins by fishermen (Northridge, 1984; UNEP/IUCN, 1994; Fertl & Leatherwood, 1997; Bearzi, 2002).

According to Bearzi *et al.* (2004), global warming is today a major concern for Mediterranean common dolphin which feed on species that are targets of fisheries, such as European anchovy, European pilchard (*Sardina pilchardus*), round sardinella (*Sardinella aurita*) and sprat (*Sprattus sprattus*) (Orsi Relini & Relini, 1993; Boutiba & Abdelghani, 1995; Birkun, 2002; Bearzi *et al.*, 2003) and that could be affected by global warming. For instance, in the Black Sea, the two mass mortality events involving common dolphins (*Delphinus delphis*), in 1990 and 1994, coincided with the decline of European sprat and anchovy stocks, their main prey species (Krivokhizhin & Birkun, 1999; Birkun, 2002). The combination of several factors, including seawater eutrophication, which is likely to be encouraged with global warming, and over-fishing, were responsible for the rapid decline of sprats and anchovies (Zaitsey & Mamaev, 1997).

OTHER POSSIBLE CLIMATE CHANGE-INDUCED IMPACTS ON CETACEANS

Climate change is likely to affect cetacean populations in several other ways (Figure 2). For instance, species that breed, feed and calve in coastal areas are more likely to be impacted by climate change which is likely to generate coastal inundations and reduce water quality by releasing pollutants into the marine environment (Orr *et al.*, 1992; Agardy, 1996; Simmonds & Nunny, 2002).

Changes in current patterns could alter cetacean migration routes and affect the transmission of sound and therefore the cetaceans' communication capacity (Agardy, 1996; MacGarvin & Simmonds, 1996; IWC, 1997).

Moreover, a change in environmental parameters like salinity might trigger more physiological stress (possibly presenting as skin lesions) and make cetaceans more susceptible to diseases or anthropogenic pressures (Wilson *et al.*, 1999; Learmonth *et al.*, 2006). These pressures, including incidental capture in fishing nets, noise and chemical pollution could have synergistic or cumulative impacts with climate change.

Finally, the increased frequency of poisonous algal blooms, like dinoflagellates, which often generate brevetoxins, has been correlated with the collapse of several marine species including cetaceans like striped dolphins in the Mediterranean Sea (Geraci *et al.*, 1989; Fitzgerald, 1991; Burns, 1998, 2001, 2002; Balmer-Hanchey *et al.*, 2003; Danovaro, 2003). Global warming, which is likely to encourage this phenomenon, could indirectly, yet profoundly affect cetaceans.

CONCLUSION

This review illustrates the different linkages existing between climate and biodiversity. It illustrates the urgent necessity for more integrated regulations for the protection of marine biodiversity. Similarly, the development of exemplary and reproducible projects aiming to reduce greenhouse gas emissions should be supported.

By influencing seawater properties, climate change can alter ecological interactions between trophic levels and is likely to disrupt overall ecosystem function. Climate change is affecting and will continue to affect marine ecosystems. Although recent climatic anomalies represent only a small part of the forecasted changes, they have already generated important responses in marine ecosystems. Marine biodiversity, including cetaceans, is highly vulnerable to environmental alteration, and can be significantly and irremediably affected by even small temperature changes. Combined with other anthropogenic pressures, climate change is likely to impact the survival of some rare and endangered marine flora and fauna, and may also threaten many other species. For example, some cetacean species that inhabit restricted geographical zones, with no option to shift their range, or those unable to effectively switch prey species if necessary may be adversely affected by climate change. As well as impacts on populations of marine biota, the physiology of individual organisms may also be severely affected, either directly or indirectly, by climate change, for example through ocean acidification.

Today, the International Whaling Commission considers global warming as a major issue (IWC, 1997) and is in the process of establishing a special workshop to examine its impacts and as Baker (in Burns, 2001) said: 'While we debate the limits that should be placed on whaling in order to protect the status of the stocks, a silent menace threatens to destroy the populations we strive to protect'. Climate change must be considered as a priority and uncertainties concerning its future impacts on biodiversity depend on the social and economic response of our societies to global warming.

REFERENCES

Abad E., Valeiras J., Gómez M.J., García-Isarch E., Baro J. and Camiñas J.A. (in press) Interactions of common dolphin *Delphinus delphis* and bottlenose dolphin *Tursiops truncatus* with trawl and purse seine fisheries at Alboràn Sea (Western Mediterranean Sea). *European Research on Cetaceans* 16.

Agardy T. (1996) *Prospective climate change impacts on cetaceans and its implications for the conservation of whales and dolphins*. Paper SC/M96/CC33 presented to the IWC, 1996. (Unpublished) [Paper available from the IWC office.]

Agostini V.N. and Bakun A. (2002) "Ocean Triads" in the Mediterranean Sea: physical mechanisms potentially structuring reproductive habitat suitability (example application to European anchovy, *Engraulis encrasicolus*). *Fisheries Oceanography* 11, 129–142.

Aguilar A. (2000) Population biology, conservation threats and status of Mediterranean striped dolphins (*Stenella coeruleoalba*). *Journal of Cetacean Research and Management* 2, 17–26.

Aguilar A., Borrel A., Calzada N. and Grau E. (1991) Body fat reserves in striped dolphins examined during the western Mediterranean die-off. In Pastor X. and Simmonds M. (eds) *Proceedings of the Mediterranean Striped Dolphin Mortality International Workshop 45, November 1991, Palma de Mallorca*. Palma de Mallorca, Spain: Greenpeace, pp. 47–52.

Aguilar A. and Raga J.A. (1993) The striped dolphin epizootic in the Mediterranean Sea. *Ambio* 22, 524–528.

Aguilar A. and Borrell A. (1994) Abnormally high polychlorinated biphenyl levels in striped dolphins (*Stenella coeruleoalba*) affected by the 1990–1992 Mediterranean epizootic. *Science of the Total Environment* 154, 237–247.

Aguilar A., Borrell A. and Pastor T. (1999) Biological factors affecting variability of persistent pollutant levels in cetaceans. In Reijnders P.J.H., Aguilar A. and Donovan G.P. (eds) *Chemical pollutants and cetaceans*, Special Issue 1. Cambridge: International Whaling Commission, pp. 83–116.

Alheit J. and Hagen E. (1997) Long-term climate forcing of European herring and sardine populations. *Fisheries Oceanography* 6, 130–139.

Álvarez M., Perez F.F., Shoosmith D.R. and Bryden H.L. (2005) Unaccounted role of Mediterranean Water in the drawdown of anthropogenic carbon. *Journal of Geophysical Research* 110, C09S03, doi:10.1029/2004JC002633.

Anadón R., Danovaro R., Dippner J.W., Drinkwater K.F., Hawkins S.J., O'Sullivan G., Oguz T. and Reid P.C. (2007) Climate change impacts on European marine and coastal environments. Ecosystems approach. *European Science Foundation Position Paper 9. Marine Board*.

Archer S., Hirano J. and Vallerga S. (2001) The effects of variability of marine ecosystems on the visual sensitivity of fish. *Archivio di Oceanografia e Limnologia* 22, 119–126.

Arkhipkin A.I., Grzebielec R., Sirota A.M., Remeslo A.V., Polishchuk I.A. and Middleton D.A.J. (2004) The influence of seasonal environmental changes on the ontogenetic migrations of the squid *Loligo gahi* on the Falkland shelf. *Fisheries Oceanography* 13, 1–9.

Astraldi M., Bianchi C.N., Gasparini G.P. and Morri C. (1995) Climatic fluctuations, current variability and marine species distribution: a case study in the Ligurian Sea (north-west Mediterranean). *Oceanologica Acta* 18, 139–149.

Astraldi M., Gasparini G.P., Ribera D'Alcala M., Conversano F. and Lavezza R. (2002) Hydrographic and chemical time series in the central Mediterranean: a sensitivity test for long-term changes in the Mediterranean Sea. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*. pp. 47–50. [Available at: http://www.ciesm.org/publications/Monaco02.pdf. Accessed 7 June 2007.]

Astruc G. and Beaubrun P. (2001) Fin whale (*Balaenoptera physalus*) summer feeding in the north-western Mediterranean Sea. In *Proceedings of the 15th European Cetacean Society Conference, Roma, 6–10 May 2001* 15, 289–291.

Athanassios C.T. and Antonopoulou E. (2006) Reproductive biology of round sardinella (*Sardinella aurita*) in the north-eastern Mediterranean. *Scientia Marina* 70, 281–290.

Azzali M., De Felice A., Cosimi G., Luna M. and Parmiggiani F. (2002) The state of the Adriatic Sea centered on the small pelagic fish populations. *PSZN Marine Ecology* 23, 78–91.

Balmer-Hanchey E.L., Jaykus L.A., Green D.P. and McClellan-Green P. (2003) Marine biotoxins of algal origin and seafood safety. *Journal of Aquatic Food Product Technology* 1, 30–53.

Bannister J.L. (2002) Baleen whales. In Perrin W.F., Würsig B. and Thewissen J.G.M. (eds) *Encyclopedia of marine mammals*. San Diego: Academic Press, pp. 62–72.

Basilone G., Guisande C., Patti B., Mazzola S., Cuttitta A., Bonanno A., Vergara A.R. and Maneiro I. (2006) Effect of habitat conditions on reproduction of the European anchovy (*Engraulis encrasicolus*) in the Strait of Sicily. *Fisheries Oceanography* 15, 271–280.

Bass C.L., Simmonds M.P., Isaac S.J. (2006) An overview of the potential consequences for cetaceans of oceanic acidification. Paper presented to the Scientific Committee of the International Whaling Commission, SC/58/E10.

Batten S. (2007) Changes in the oceanic northeast Pacific plankton populations; what may happen in a warmer ocean. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 2 June 2007.]

Beare D.J., Batten S., Edwards M. and Reid D.G. (2002) Prevalence of boreal Atlantic, temperate Atlantic and neritic zooplankton in the North Sea between 1958 and 1998 in relation to temperature, salinity, stratification intensity and Atlantic inflow. *Journal of Sea Research* 48, 29–49.

Bearzi G., Politi E. and Notarbartolo di Sciara G. (1999) Diurnal behavior of free-ranging bottlenose dolphins in the Kvarneric (northern Adriatic Sea). *Marine Mammal Science* 15, 1065–1097.

Bearzi G. (2002) Interactions between cetaceans and fisheries in the Mediterranean Sea. In Notarbartolo di Sciara G. (ed.) *Cetaceans of the Mediterranean and Black Seas: state of knowledge and conservation strategies*. A report to the ACCOBAMS Secretariat, Monaco, February 2002. Section 9, 20 p. [Available at http://www.accobams.org/download/ceta_mbs_report/09_bearzi.pdf; accessed 2 June 2007.]

Bearzi G., Reeves R.R., Notarbartolo di Sciara G., Politi E., Cañadas A., Frantzis A. and Mussi B. (2003) Ecology, status and conservation of short-beaked common dolphins (*Delphinus delphis*) in the Mediterranean Sea. *Mammal Review* 33, 224–252.

Bearzi G., Notarbartolo di Sciara G., Reeves R.R., Canadas A. and Frantzis A. (2004) *Conservation Plan for short-beaked common dolphins in the Mediterranean Sea. ACCOBAMS, Agreement on the Conservation of Cetaceans of the Black Sea, Mediterranean Sea and Contiguous Atlantic Area*, 90 pp.

Beaugrand G. (2003) Long-term changes in copepod abundance and diversity in the north-east Atlantic in relation to fluctuations in the hydroclimatic environment. *Fisheries Oceanography* 12, 270–283.

Beaugrand G. and Ibanez F. (2004) Monitoring marine plankton ecosystems II: long-term changes in North Sea calanoid copepods in relation to hydro-climatic variability. *Marine Ecology Progress Series* 284, 35–47.

Beaugrand G. and Reid P.C. (2003) Long-term changes in phytoplankton, zooplankton and salmon related to climate. *Global Change Biology* 9, 801–817.

SC/N10/CC For Info 2

Beaugrand G., Reid P.C., Ibanez F., Lindley J.A. and Edwards M. (2002) Reorganization of North Atlantic marine copepod biodiversity and climate. *Science* 296, 1692–1694.

Beaugrand G., Brander K.M., Lindley J.A., Souissi S. and Reid P.C. (2003) Plankton effect on cod recruitment in the North Sea. *Nature* 426, 661–664.

Besson J., Duguy R. and Tardy G. (1982) Note sur un cas de multiparité chez un rorqual commun (*Balaenoptera physalus*). *Mammalia* 46, 408.

Béthoux J.P. and Gentili B. (1996) The Mediterranean Sea, coastal and deep-sea signatures of climatic and environmental changes. *Journal of Marine Systems* 7, 383–394.

Béthoux J.P., Gentili B., Raunet J. and Tailliez D. (1990) Warming trend in the western Mediterranean deep water. *Nature* 347, 660–662.

Béthoux J.P., Gentili B. and Tailliez D. (1998) Warming and freshwater budget change in the Mediterranean since the 1940s, their possible relation to the greenhouse effect. *Geophysical Research Letters* 25, 1023–1026.

Béthoux J.P., Morin P. and Ruiz-Pino D.P. (2002) Temporal trends in nutrient ratios: chemical evidence of Mediterranean ecosystem changes driven by human activity. *Deep-Sea Research, II* 49, 2007–2016.

Béthoux J.P., Gentili B., Morin P., Nicolas E., Pierre C. and Ruiz-Pino D. (1999) The Mediterranean Sea: a miniature ocean for climatic and environmental studies and a key for the climatic functioning of the North Atlantic. *Progress in Oceanography* 44, 131–146.

Bianchi C.N. (1997) Climate change and biological response in the marine benthos. *Atti Associazione Italiana Oceanologia Limnologia* 1, 3–20.

Bianchi C.N. (2007) Biodiversity issues for the forthcoming tropical Mediterranean Sea. *Hydrobiologia* 580, 7–21.

Bianchi C.N. and Morri C. (1993) Range extensions of warm-water species in the northern Mediterranean: evidence for climatic fluctuations? *Porcupine Newsletter* 5, 156–159.

Bianchi C.N. and Morri C. (1994) Southern species in the Ligurian Sea (northern Mediterranean): new records and a review. *Bollettino dei Musei e Degli Istituti Biologici dell'Università di Genova* 58–59, 181–197.

Bianchi C.N. and Morri C. (2000) Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research. *Marine Pollution Bulletin* 40, 367–376.

Birkun A.J. (2002) Interaction between cetaceans and fisheries in the Black Sea. In Notarbartolo di Sciara G. (ed.) *Cetaceans of the Mediterranean and Black Seas: state of knowledge and conservation strategies*. A report to the ACCOBAMS Secretariat, Monaco, February 2002, pp. 98–107.

Bjorge A. (2002) How persistent are marine mammal habitats in an ocean of variability? In Evans P.G.H. and Raga J.A. (eds) *Marine mammals: biology and conservation*. New York: Kluwer Academic/Plenum Publishers, pp.63–91.

Blanco C., Salomon O. and Raga J.A. (2001) Diet of the bottlenose dolphin (*Tursiops truncatus*) in the western Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom* 81, 1053–1058.

Blunier T. (2000) Frozen methane escapes from the sea floor. *Science* 288, 68–69.

Boero F. (1996) Episodic events: their relevance to ecology and evolution. *Marine Ecology*, 17, 237–250.

Bombace G. (2001) Influence of climatic changes on stocks, fish species and marine ecosystems in the Mediterranean Sea. *Archives of Oceanography and Limnology* 22, 67–72.

Bompar J.M. (2000) *Les cétacés de Méditerranée*. Edisud, La Calade: Aix-en-Provence. 188 pp.

Bopp L., Legendre L. and Monfray P. (2002) La pompe à carbone va-t-elle se gripper? *La Recherche (spécial La Mer)* 355, 48–51.

Bouderesque C.F. and Verlaque M. (2002) Biological pollution in the Mediterranean Sea: invasive versus introduced macrophytes. *Marine Pollution Bulletin* 44, 32–38.

Boutiba Z. and Abdelghani F. (1995) Food of the common dolphin (*Delphinus delphis*, L.) in Algerian waters. *European Research on Cetaceans* 9, 182.

Bowen W.D. (1997) The role of marine mammals in aquatic ecosystems. *Marine Ecology Progress Series* 158, 267–274.

Boyd I.L. (2002) Antarctic marine mammals. In Perrin W.F., Würsig B. and Thewissen J.G.M. (eds) *Encyclopedia of marine mammals*. San Diego: Academic Press, pp. 30–36.

Bräger S. (1993) Diurnal and seasonal behavior patterns of bottlenose dolphins (*Tursiops truncatus*). *Marine Mammal Science* 9, 434–437.

Brander K. (2005) Cod recruitment is strongly affected by climate when stock biomass is low. *ICES Journal of Marine Science* 62, 339–343.

Brodeur R.D., Mills C.E., Overland J.E., Walters G.E. and Schumacher J.D. (1999) Evidence for a substantial increase in gelatinous zooplankton in the Bering Sea, with possible links to climate change. *Fisheries Oceanography* 8, 296–306.

Buchholz F., Buchholz C., Reppin J. and Fisher J. (1995) Diel vertical migrations of *Meganyctiphanes norvegica* in the Kattegat: comparison of net catches and measurements with Acoustic Doppler Current Profilers. *Helgoländer Wissenschaftliche Meeresuntersuchungen* 49, 849–866.

Burns W.C.G. (1998) 'The Agreement on the Conservation of Cetaceans of the Black Sea, Mediterranean Sea and Contiguous Atlantic Area: The Regional Response to the Threats Facing Cetaceans.' *Journal of International Wildlife Law and Policy* 1, 113–133.

Burns W.C.G. (2001) From the harpoon to the heat: climate change and the International Whaling Commission in the 21st century. *Georgetown International Environmental Law Review* 13, 335–359. [Available at: http://www.pacinst.org/press_center/IWCOP.pdf; accessed 5 July 2007.]

Burns W.C.G. (2002) Climate change and the International Whaling Commission in the 21st century. In Burns W.G.C. and Gillespie A. (eds) *The future of cetaceans in a changing world*. New York: Transnational Publishers, pp. 339–379.

Calbet A., Broglio E., Saiz E. and Alcaraz M. (2002) Low grazing impact of mesozooplankton on the microbial communities of the Alborán Sea: a possible case of inhibitory effects by the toxic dinoflagellate *Gymnodinium catenatum*. *Aquatic Microbiology and Ecology* 26, 235–246.

Caldeira K. and Wickett M.E. (2003) Anthropogenic carbon and ocean pH. *Nature* 425, 365.

Caldwell D.K. and Caldwell M.C. (1978) Cetaceans. In Layne J.N. (ed.) *Rare and endangered biota of Florida, Volume 1: mammals*. Gainesville, FL: University of Florida Press, pp. 49–52.

Carlton J.T. (2000) Global change and biological invasions in the oceans. In Mooney H.A. and Hobbs R.J. (eds) *Invasive species in a changing world*. Covelo, CA: Island Press, pp. 31–53.

CBD (2003) *Secretariat of the Convention on Biological Diversity. Interlinkages between biological diversity and climate change. Advice on the integration of biodiversity considerations into the implementation of the United Nations Framework Convention on Climate Change and its Kyoto Protocol*. Montreal, SCBD, 154

pp. (CBD Technical Series no. 10). [Available at: http://www.cbd.int/doc/publications/cbd-ts-10.pdf; accessed 15 July 2007.]

Cebrian D. (1995) The striped dolphin *Stenella coeruleoalba* epizootic in Greece, 1991–1992. *Biological Conservation* 74, 143–145.

Cerrano C., Bavestrello G., Bianchi C.N., Cattaneo-Vietti R., Bava S., Morganti C., Morri C., Picco P., Sara G., Schiaparelli S., Siccardi A. and Sponga F. (2000) A catastrophic mass-mortality episode of gorgonians and other organisms in the Ligurian Sea (North-western Mediterranean), summer 1999. *Ecology Letters* 3, 284–293.

Cerrano C., Totti C., Sponga F. and Bavestrello G. (2006) Summer disease in *Parazoanthus axinellae* (Schmidt, 1862) (Cnidaria, Zoanthidea). *Italian Journal of Zoology* 73, 355–361.

Chevaldonné P. and Lejeusne C. (2003) Regional warming induced species shift in north-west Mediterranean marines caves. *Ecology Letters* 6, 371–379.

Cheviron N. (2007) Le réchauffement climatique menace la Méditerranée. www.cyberpress.ca [Accessed 13 April 2007.]

CIESM (2000) *Fishing down the Mediterranean food web?* CIESM Workshop Series. Kerkyra, Greece, 26–30 July 2000. 99 pp.

CIESM (2004) Novel contaminants and pathogens in coastal waters. *CIESM Workshop Monograph* 26, Monaco, 116 pp.

CIESM (2005) *Atlas of exotic species in the Mediterranean*. [Available at http://www.ciesm.org/atlas; accessed 7 July 2007.]

Clark R.A., Fox C.J., Viner D. and Livermore M. (2003) North Sea cod and climate change—modelling the effects of temperature on population dynamics. *Global Change Biology* 9, 1669–1680.

Cohen J. and Barlow M. (2005) The NAO, the AO, and global warming: how closely related? *Journal of Climate* 18, 4498–4513.

COSEWIC (2005) *Assessment and update status report on the bowhead whale* Balaena mysticetus *in Canada*. Committee on the Status of Endangered Wildlife in Canada. Ottawa. viii + 51 pp. [Available at: http://www.sararegistry.gc.ca/virtual_sara/files/cosewic/sr%5Fbowhead%5Fwhale%5Fe%2Epdf; accessed 4 July 2007.]

Cotton P.A., Sims D.W., Fanshawe S. and Chadwick M. (2005) The effects of climate variability on zooplankton and basking shark relative abundance off southwest Britain. *Fisheries Oceanography* 14, 151–155.

Cox P., Betts R., Jones C., Spall S. and Totterdell I. (2000) Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature* 408, 184–187.

Cury P. and Roy C. (1989) Optimal environmental window and pelagic fish recruitment success in upwelling areas. *Canadian Journal of Fisheries and Aquatic Science* 46, 670–680.

Danovaro R. (2003) Pollution threats in the Mediterranean Sea: an overview. *Chemistry and Ecology* 19, 15–32.

Danovaro R., Dell'Anno A., Fabiano M. and Pusceddu A. (2004) Biodiversity response to climate change in a warm deep sea. *Ecology Letters* 7, 821–828.

Danovaro R., Dell'Anno A., Fabiano M., Pusceddu A. and Tselepides A. (2001) Deep-sea ecosystem response to climate changes: the eastern Mediterranean case study. *Tree* 16, 505–510.

Danovaro R., Dell'Anno A., Fabiano M., Pusceddu A. and Tselepides A. (2001) Deep-sea ecosystem response to climate changes: the eastern Mediterranean case study. *Trends in Ecology and Evolution* 16, 505–510.

Degobbis D., Precali R., Ivancic I., Smodlaka N., Fuks D. and Kveder S. (2000) Long-term changes in the northern Adriatic ecosystem related to anthropogenic eutrophication. *International Journal of Environment and Pollution* 13, 495–553.

De Heij A. and Baayen R.P. (2005) Seasonal distribution of cephalopod species living in the central and southern North Sea. *Basteria* 69, 91–119.

De la Bigne C. (1985) *Etude du métabolisme nutritionnel et des variations spatio-temporelles de l'environnement trophique potentiel de Meganyctiphanes norvegica (Euphausiacea) en Mer Ligure*. Thèse de Doctorat en océanographie biologique. Université Pierre et Marie Curie (Paris VI), 137 pp.

Dhermain F. (2003) Suivi Hivernal et Estival de Grand Dauphin (*Tursiops truncatus*) en Corse en 2002 et 2003. *12ème Conférence Internationale RI.MO*. Palais des Congrès D'Antibes-Juan-les Pins 15 November 2003.

Dhermain F., Soulier L. and Bompar J.M. (2002) Natural mortality factors affecting cetaceans in the Mediterranean Sea. In G. Notarbartolo di Sciara (ed.) *Cetaceans of the Mediterranean and Black Seas: state of knowledge and conservation strategies. A report to the ACCOBAMS Secretariat, Monaco, February 2002. Section 15*, 14 pp.

Drake L.A., Doblin M.A. and Dobbs F.C. (2007) Potential microbial bioinvasions via ships' ballast water, sediment, and biofilm. *Marine Pollution Bulletin* 55, 333–341.

Duarte C.M., Agustí S., Kennedy H. and Vaqué D. (1999) The Mediterranean climate as a template for Mediterranean marine ecosystems: the example of the northeast Spanish littoral. *Progress in Oceanography* 44, 245–270.

Ducrotoy J.P. (1999) Indications of change in the marine flora of the North Sea in the 1990s. *Marine Pollution Bulletin* 38, 646–654.

Duffy J.E. (2003) Biodiversity loss, trophic skew and ecosystem functioning. *Ecology Letters* 6, 680–687.

Dulcic J. and Grbec B. (2000) Climate change and Adriatic ichthyofauna. *Fisheries Oceanography* 9, 187–191.

Dulcic J., Grbec B. and Lipej L. (1999) Information on the Adriatic ichthyofauna—effect of water warming? *Acta Adriatica* 40, 33–43.

Dulcic J. and Lipej L. (1997) New records of marine fishes from the Slovenian coastal waters. *Falco* 12, 35–40.

Dulcic J., Tutman P. and Caleta M. (2006) Northermost occurrence of the white grouper, *Epinephelus aeneus* (Perciformes: Serranidea), in the Mediterranean area. *Acta Ichthyologica et Piscatoria* 36, 73–75.

Dulcic J., Kraljevic M., Pallaoro A. and Glamuzina B. (2005) Unusual catch of bluefish *Pomatomus saltatrix* (Pomatomidae) in Tarska cove (northern Adriatic). *Cybium* 29, 207–208.

Edwards M. and Beaugrand G. (2007) Spatial variability of marine climate change impacts in the North Atlantic. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 2 of June 2007.]

Edwards M. and Richardson A.J. (2004) Impact of climate change on marine pelagic phenology and trophic mismatch. *Nature* 430, 881–884.

Edwards M., Beaugrand G., Reid P.C., Rowden A.A. and Jones M.B. (2002) Ocean climate anomalies and the ecology of the North Sea. *Marine Ecology Progress Series* 239, 1–10.

Edwards M., Reid P. and Planque B. (2001) Long-term and regional variability of phytoplankton biomass in the Northeast Atlantic (1960–1995). *ICES Journal of Marine Science* 58, 39–49.

EEA (2004) *Impacts of Europe's changing climate. An indicator-based assessment*. Copenhagen: EEA.

SC/N10/CC For Info 2

Einarsson H. (1945) Euphausiacea. 1. North Atlantic species. *Dana Report* 27, 1–185.

Elliot M. (2003) Biological pollutants and biological pollution—an increasing cause for concern. *Marine Pollution Bulletin* 46, 275–280.

Estrada M. (1996) Primary production in the northwestern Mediterranean. *Scientia Marina* 60, 55–64.

European Community (1999) State and pressures of the marine and coastal Mediterranean environment. *Environmental Issues Series* 5, 137.

Evans W.E. (1971) Orientation behavior of delphinids: radio telemetric studies. *Annals of the New York Academy of Sciences* 188, 142–160.

Feely R., Sabine C., Lee K., Berelson W., Kleypas J., Fabry V. and Millero F. (2004) Impact of anthropogenic CO_2 on the $CaCO_3$ system in the oceans. *Science* 305, 362–366.

Fernandez de Puelles M.L., Valencia J. and Vicente L. (2004) Zooplankton variability and climatic anomalies from 1994 to 2001 in the Balearic Sea (Western Mediterranean). *ICES Journal of Marine Science* 61, 492–500.

Fertl D. and Leatherwood S. (1997) Cetacean interactions with trawls: a preliminary review. *Journal of Northwest Atlantic Fishery Science* 22, 219–248.

Fisher S., Hanley K., Lonsdale J., Nelson K. and Thornton A. (1994) *Under fire: environmental threats and extinction of the world's cetaceans*. London and Washington, DC: Environmental Investigation Agency, 33 pp.

Fitzgerald J.W. (1991) Marine aerosols: a review. *Atmospheric Environment* 25A, 533–545.

Font J., Puig P., Salat J., Palanques A. and Emelianov M. (2007) Sequence of hydrographic changes in the NW Mediterranean deep water due to the exceptional winter 2005. *Scientia Marina* 71, 339–346.

Forcada J., Aguilar A., Hammond P., Pastor X. and Aguilar R. (1996) Distribution and abundance of fin whales (*Balaenoptera physalus*) in the Western Mediterranean Sea during the summer. *Journal of Zoology, London* 238, 23–34.

Forward R.B. and Fyhn H.J. (1983) Osmotic regulation of the krill *Meganyctiphanes norvegica*. *Comparative Biochemistry and Physiology* 74, 301–305.

Foster B.A. (1971) On the determinants of the upper limit of intertidal distribution of barnacles (Crustacea: Cirripedia). *Journal of Animimal Ecology* 40, 33–48.

Fowler S.W., Small L.F. and Keckes S. (1971) Effects of temperature and size on molting of euphausiid crustaceans. *Marine Biology* 11, 45–51.

Francour P. (1997) Fish assemblages of *Posidonia oceanica* beds at Port-Cros (France, NW Mediterranean): assessment of composition and long-term fluctuations by visual census. *Marine Ecology* 18, 157–173.

Francour P., Boudouresque C.F., Harmelin J.G., Harmelin-Vivien M.L. and Quignard J.P. (1994) Are the Mediterranean waters becoming warmer? Information from biological indicators. *Marine Pollution Bulletin* 28, 523–526.

Frank K.T., Petrie B., Choi J.S. and Leggete W.C. (2005) Trophic cascades in a formerly cod-dominated ecosystem. *Science* 308, 1621–1623.

Fromentin J.M. and Planque B. (1996) *Calanus* and environment in the eastern North Atlantic. II. Influence of the North Atlantic Oscillation on *C. finmarchicus* and *C. helgolandicus*. *Marine Ecology Progress Series* 134, 111–118.

Fuda J.L. and Millot C. (2002) Warming and salting of the Tyrrhenian Deep Water. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series 16, 22–24 April, Monaco*, pp. 51–54. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Fullard K.J., Early G., Heide-Jorgensen M.P., Bloch D., Rosing-Avid A. and Amos W. (2000) Population structure of long-finned pilot whales in the North Atlantic: a correlation with sea surface temperature? *Molecular Ecology* 9, 949–958.

Fuller G.B. and Hobson W.C. (1986) Effects of PCBs on reproduction in mammals. In Waid J.S. (ed.) *PCBs and the environment*, Volume II. Boca Raton, FL: CRC Press, pp. 101–125.

Gacic M. (2002) Southern Adriatic and Otranto Strait—key areas for climatic monitoring. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 55–56. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Galil B.S. and Zenetos A. (2002) A sea change—exotics in the eastern Mediterranean. In Leppakoski E., Gollasch S. and Olenin S. (eds.) *Invasive aquatic species of Europe. Distribution, impacts and management*. Dordrecht: Kluwer Academic Publishers, pp. 325–336. [Available at: http://www.biomareweb.org/downloads/exotic_med.pdf; accessed 10 June 2007.]

Galil B.S. (2007) Loss or gain? Invasive aliens and biodiversity in the Mediterranean Sea. *Marine Pollution Bulletin* 55, 314–322.

Galil B.S., Nehring S. and Panov V. (2007) Waterways as invasion highways—impact of climate change and globalization. In Nentwig W. (ed.) *Biological invasions*. (Ecological Studies, 193). Berlin and Heidelberg: Springer, pp. 59–74.

Gambaiani D., Mayol P. and Isaac S. (2005) Potential impacts of global warming on cetaceans. *20th ECS Conference*, 2–7 April 2006, Gdynia, Poland.

Gannier A. (1995) *Les Cétacés de Méditerranée Nord-Occidentale: estimation de leur abondance et mise en relation de la variation saisonnière d leur distribution avec l'écologie du milieu*. Thèse de doctorat. Ecole Pratiques des Hautes Etudes, Montpellier, France. 433 pp.

Gannier A. (1997) Estimation de l'abondance estivale du rorqual commun *Balaenoptera physalus* (Linné, 1758) dans le bassin Ligure-Provençal (Méditerranée occidentale). *Revue Ecologie (Terre Vie)* 52, 69–86.

Gantzer C., Dubois E., Crance J.M., Billaudel S., Kopecka H., Schwartzbrod L., Pommepuy M. and Le Guyader F. (1998) Devenir des virus entériques en mer et influence des facteurs environnementaux. *Oceanologica Acta* 21, 983–992.

Garcés E., Maso M., Vila M. and Camp J. (2000) Harmful algae events in the Mediterranean: are they increasing? *Harmful Algae News* 20, 10–11.

Garcia A. and Palomera I. (1996) Anchovy early life history and its relation to its surrounding environment in the Western Mediterranean basin. *Scientia Marina* 60 (Supplement 2), 155–166.

Garrabou J., Perez T., Sartoretto S. and Harmelin J.G. (2001) Mass mortality event in red coral *Corallium rubrum* populations in the Provence region (France, NW Mediterranean). *Marine Ecology Progress Series* 217, 263–272.

Gaskin D.E. (1968) Distribution of Delphinidea (Cetacea) in relation to sea surface temperature off eastern and southern New Zealand. *New Zealand Journal of Marine and Freshwater Research* 2, 527–534.

Gasparini G.P. and Astraldi M. (2002) Experimental evidence of the interannual variability of currents in two Mediterranean straits: the Strait of Sicily and the Corsica Channel. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16,*

SC/N10/CC For Info 2

22–24 April, Monaco, pp. 43–46. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Genner M.J., Sims D.W., Wearmouth V.J., Southall E.J., Southward A.J., Henderson P.A. and Hawkins S.J. (2004) Regional climatic warming drives long-term community changes of British marine fish. *Proceedings of the Royal Society of London Series B: Biological Sciences* 271, 655–661.

Geraci J.R. and Lounsbury V. (2002) Marine mammal health: holding the balance in an ever changing sea. In Evans P.G.H and Raga J.A. (eds) *Marine mammals: biology and conservation*. New York: Kluwer Academic/Plenum Publishers, pp. 365–384.

Geraci J.R., Andersen D.M., Timperi R.J., St Aubin D.J., Early G.A., Prescott J.H. and Mayo C.A. (1989) Humpback whales fatally poisoned by dinoflagellate toxin. *Canadian Journal of Fisheries and Aquatic Sciences* 46, 1895–1898.

Gertman I. and Hecht A. (2002) Annual and long-term changes in the salinity and the temperature of the waters of the South-eastern Levantine Basin. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 75–78. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Giannakopoulos C., Bindi M., Moriondo M., LeSager P. and Tin T. (2005) *Climate change impacts in the Mediterranean resulting from a 2°C global temperature rise*, WWF Report, pp. 66. [Available at: http://assets.panda.org/downloads/medreportfinal8july05.pdf; accessed 2 June 2007.]

Gillet N.P., Graf H.F and Osborn T.J. (2003) *Climate change and the NAO*. Washington, DC: American Geophysical Union.

Giorgi F. (2006) Climate change hot-spots. *Geophysical Research Letters* 33, L08707.

Glamuzina B. (1999) Recent changes of Adriatic ichthyofauna: threat or benefit to local fishery. *CIESM Workshop Series* 7, 71–73.

Goffart A., Hecq J.H. and Legendre L. (2002) Changes in the development of the winter–spring phytoplankton bloom in the bay of Calvi (NW Mediterranean) over the last two decades: a response to the changing climate? *Marine Ecology Progress Series* 236, 45–60.

Gomez F. (2003) The toxic dinoflagellate *Gymnodinium catenatum*: an invader in the Mediterranean Sea. *Acta Botanica Croatica* 62, 65–72.

Gomez F. and Claustre H. (2003) The genus *Asterodinium* (Dinophyceae) as a possible biological indicator of warming in the western Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom* 83, 173–174.

Graham N.E. (1995) Simulation of recent global temperature trends. *Science* 267, 666–671.

Grebmeier J.M. and Dunton K.H. (2000) Benthic processes in the northern Bering/Chukchi Seas: status and global change. In Huntington H.P. (ed.) *Impacts of changes in sea ice and other environmental parameters in the Arctic. Report of the Marine Mammal Commission Workshop, 15–17 February 2000*. Bethesda: Marine Mammal Commission, pp. 61–71.

Green E.J. and Carritt D.E. (1967) Oxygen solubility in sea water: thermodynamic influence of sea salt. *Science* 157, 191–193.

Greene C.H. and Pershing A.J. (2004) Climate and the conservation biology of North Atlantic right whales: the right whale at the wrong time? *Frontiers in Ecology and the Environment* 2, 29–34.

Greve W. (2007) Climate effects on North Sea zooplankton. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 2 June 2007.]

Gritti E.S., Smith B. and Sykes M.T. (2006) Vulnerability of Mediterranean basin ecosystems to climate change and invasion by exotic plant species. *Journal of Biogeography* 33, 145–157.

Guidetti P., Boero F. and Dulcic J. (2002) Mass mortality of gilt sardine, *Sardinella aurita* (Clupeidae), in the Adriatic and Ionian Seas. *Cybium* 26, 317–319.

Guidetti P. and Boero F. (2001) Occurrence of the Mediterranean parrotfish *Sparisoma cretense* (Perciformes: Scaridae) in south-eastern Apulia (south-east Italy). *Journal of the Marine Biological Association of the United Kingdom* 81, 717–718.

Gulland F.M.D., Perez-Cortes M., Urban R., Rojas-Bracho L., Ylitalo G., Weir J., Norman S.A., Muto M.M., Rugh D.J., Kreuder C. and Rowles T. (2005) *Eastern North Pacific gray whale* (Eschrichtius robustus) *unusual mortality event, 1999–2000*. NOAA Technical Memorandum NMFS-AFSC-150-150. US Department of Commerce, Washington, DC, USA.

Hannah L., Midgley G.F., Lovejoy T., Bond W.J., Bush M., Lovett J.C., Scott D. and Woodward F.I. (2002) Conservation of biodiversity in a changing climate. *Conservation Biology* 16, 264–268.

Hanson M.T. and Defran R.H. (1993) The behaviour and feeding ecology of the Pacific coast bottlenose dolphin, *Tursiops truncatus*. *Aquatic Mammals* 19, 127–142.

Hardwood J. (2001) Marine mammals and their environment in the twenty-first century. *Journal of Mammology* 82, 630–640.

Harley C.D.G., Hughes A.R., Hultgren K.M., Miner B.G., Sorte C.J.B., Thornber C.S., Rodriguez L.F., Tomanek L. and Williams S.L. (2006) The impacts of climate change in coastal marine systems. *Ecology Letters* 9, 228–241.

Harmelin-Vivien M.L., Bitar G., Harmelin J.-G. and Monestiez P. (2005) The littoral fish community of the Lebanese rocky coast (eastern Mediterranean Sea) with emphasis on Red Sea immigrants. *Biological Invasions* 7, 625–637.

Harris R.C. Bodaly R.A. (1998) Temperature, growth and dietary effects on fish mercury dynamics in two Ontario lakes. *Biogeochemistry* 40, 175–187.

Harris L.G. and Tyrrell M.C. (2001) Changing community states in the Gulf of Maine: synergism between invaders, overfishing and climate change. *Biological Invasions* 3, 9–21.

Harvell C.D., Kim K., Burkholder J.M., Colwell R.R., Epstein P.R., Grimes D.J., Hofmann E.E., Lipp E.K., Osterhaus A.D.M.E., Overstreet R.M., Porter J.W., Smith G.W. and Vasta G.R. (1999) Emerging marine diseases—climate links and anthropogenic factors. *Science* 285, 1503–1510.

Harvell C.D., Mitchell C.E., Ward J.R., Altizer S., Dobson A.P., Ostfeld R.S. and Samuel M.D. (2002) Climate warming and disease risks for terrestrial and marine biota. *Science* 296, 2158–2162.

Heath M.R., Beare D.J., Dunn J., Fraser J.G., Hay S.J. and Turrell W.R. (1999) Monitoring the effects of climate change: overwintering abundance of *Calanus finmarchicus* in the Faroe–Shetland Channel. *Fisheries Research Services Report No. 14/99*, 24 pp.

Hernández M., Robinson I., Aguilar A., González L.M., López-Jurado L.F., Reyero M.I., Cacho E., Franco J., López-Rodas V. and Costas E. (1998) Did algal toxins cause monk seal mortality? *Nature* 393, 28.

Hesselbo S.P., Grocke D.R., Jenkyns H.C., Bjerrum C.J., Farrimond P., Morgans Bell H.S. and Green O.R. (2000) Massive dissociation of gas hydrate during a Jurassic oceanic anoxic event. *Nature* 406, 392–395.

Hiscock K., Southward A., Tittley I. and Hawkins S. (2004) Effects of changing temperature on benthic marine life in Britain and Ireland. *Aquatic Conservation: Marine and Freshwater Ecosystems* 14, 333–362.

SC/N10/CC For Info 2

Hochachka P.W. and Somero G.N. (2002) *Biochemical adaptation: mechanism and process in physiological evolution*. New York: Oxford University Press.

Hughes L. (2000) Biological consequences of global warming: is the signal already apparent? *Tree* 15, 56–61.

Hurrell J.W. (1995) Decadal trends in the North Atlantic Oscillation: regional temperature and precipitation. *Science* 269, 676–679.

Hurrell J.W. (1996) Influence of variations in extratropical wintertime teleconnections on northern hemisphere temperature. *Geophysical Research Letters* 23, 665–668.

IPCC (2001a) *Climate Change 2001: impacts, adaptation and vulnerability, a contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change (IPCC). Intergovernmental Panel on Climate Change, 3rd Assessment Report*. Cambridge: Cambridge University Press. 1000 pp.

IPCC (2001b) *Climate Change 2001: synthesis report, a contribution of Working Groups I, II and III to the Third Assessment Report of the Intergovernmental Panel on Climate Change. Intergovernmental Panel on Climate Change, 3rd Assessment Report*. Cambridge: Cambridge University Press. 398 pp.

IPCC (2007a) *Climate Change 2007: the physical science basis. Summary for policy makers. Intergovernmental Panel on Climate Change 4th Assessment Report.* Cambridge: Cambridge University Press.

IPCC (2007b) *Climate Change 2007: impacts, adaptation and vulnerability. Summary for policymakers. Intergovernmental Panel on Climate Change 4th Assessment Report.* Cambridge: Cambridge University Press.

IWC (1997) Report of the IWC workshop on climate change and cetaceans. *Report of the International Whaling Commission* 47, 293–313.

Jancovici J.M. (2005) *Changement climatique ou choc climatique?* [Available at: http://www.manicore.com; accessed 1 March 2005.]

Jefferson T.A. and Schiro A.J. (1997) Distribution of cetaceans in the offshore Gulf of Mexico. *Mammal Review* 27, 27–50.

Jones E.G., Collins M.A., Bagley P.M., Addison S. and Priede I.G. (1998) The fate of cetacean carcasses in the deep sea: observations on consumption rates and succession of scavenging species in the abyssal Northeast Atlantic Ocean. *Proceedings of the Royal Society of London Series B, Biological Sciences* 265, 1119–1127.

Justic D. (1987) Long-term eutrophication of the northern Adriatic Sea. *Marine Pollution Bulletin* 18, 281–284.

Kamburska L. and Fonda-Umani S. (2006) Long-term copepod dynamics in the Gulf of Trieste (Northern Adriatic Sea): recent changes and trends. *Climate Research* 31, 195–203.

Katona S. and Whitehead H. (1988) Are Cetacea ecologically important? *Oceanography and Marine Biology: an Annual Review* 26, 553–568.

Kennedy S. (1999) Morbilliviral infections in marine mammals. In Reijnders P.J.H., Aguilar A. and Donovan G.P. (eds) *Chemical pollutants and cetaceans*, Special Issue 1. Cambridge: International Whaling Commission, pp. 267–273.

Kennedy T.A., Naemm S., Howe K.M., Knops J.M.H., Tilman D. and Reich P. (2002) Biodiversity as a barrier to ecological invasion. *Nature* 417, 636–638.

Kennett J.P., Cannariato K.G., Hendy I.L. and Behl R.J. (2000) Carbon isotopic evidence for methane hydrate instability during Quaternary interstadials. *Science* 288, 128–133.

Kenney V.S. (1990) Anticipated effects of climate change on estuarine and coastal fisheries. *Fisheries* 15, 16–22.

Kenney R.D., Payne P.M., Heinemann D.W. and Winn H.E. (1996) Shifts in northeast shelf cetacean distributions relative to trends in Gulf of Maine/Georges Bank finfish abundance. In Sherman K., Jaworski N.A. and Smada T. (eds) *The northeast shelf ecosystem: assessment, sustainability and management*. Cambridge, MA: Blackwell Science, pp. 169–196.

Klein B., Roether W., Manca B.B., Bregant D., Beitzel V., Kovacevic V. and Luchetta A. (1999) The large deep water transient in the Eastern Mediterranean. *Deep-Sea Research* 46, 371–414.

Kleypas J.A., Buddemeier R.W., Archer D., Gattuso J.-P., Langdon C. and Opdyke B.N. (1999) Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science* 284, 118–120.

Komatsu T., Meinesz A. and Buckles D. (1997) Temperature and light responses of alga *Caulerpa taxifolia* introduced into the Mediterranean Sea. *Marine Ecology Progress Series* 146, 145–153.

Krahmann G. and Schott F. (1998) Longterm increases in Western Mediterranean salinities and temperatures: anthropogenic and climatic sources. *Geophysical Research Letters* 25, 4209–4212.

Krivokhizhin S.V. and Birkun A.J. (1999) Strandings of cetaceans along the coasts of Crimean peninsula in 1989–1996. *European Research on Cetaceans* 12, 59–62.

Kroncke I., Dippner J.W., Heyen H. and Zeiss B. (1998) Long-term changes in macrofaunal communities off Norderney (East Frisia, Germany) in relation to climate variability. *Marine Ecology Progress Series* 167, 25–36.

Kushmaro A., Rosenberg E., Fine M., Ben Haim Y. and Loya Y. (1998) Effects of temperature on the bleaching of the coral *Oculina patagonica* by the *Vibrio* AK-1. *Marine Ecology Progress Series* 171, 131–137.

Lacombe H., Tchernia P. and Gamberoni L. (1985) Variable bottom water in the Western Mediterranean basin. *Progress in Oceanography* 14, 319–338.

Lafferty K.D., Porter J.W. and Ford S.E. (2004) Are diseases increasing in the ocean? *Annual Review of Ecology and Systematics* 35, 31–54.

Lafuente J.G., Mazzola S., Quintanilla L., Delgado J., Cuttita A. and Patti B. (2002) Hydrographic phenomena influencing early life stages of the Sicilian Channel anchovy. *Fisheries Oceanography* 11, 31–44.

Lafuente J.G., Vargas J.M., Criado F., Garcia A., Delgado J. and Mazzola S. (2005) Assessing the variability of hydrographic processes influencing the life cycle of the Sicilian Channel anchovy, *Engraulis encrasicolus*, by satellite imagery. *Fisheries Oceanography* 14, 32–46.

Laidre K.L. and Heide-Jorgensen M.P. (2005) Arctic sea ice trends and narwhal vulnerability. *Biological Conservation* 121, 509–517.

Lascaratos A., Sofianos S. and Korres G. (2002) Interannual variability of atmospheric parameters over the Mediterranean basin from 1945 to 1994. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 21–23. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Laubier L. (2001) Climatic changes and trends in marine invertebrates: a need for relevant observing networks and experimental ecophysiology. *Atti Associazione Italiana Oceanologiae Limnologia* 14, 15–24.

Laubier L., Pérez T. and Garrabou J. (2003) Mass mortalities of sponges and gorgonians in the Northwest Mediterranean in 1999. *Letter PIGBPMRC France* 15, 50–56.

Le Boeuf B.J., Pérez-Cortés M.H., Urbán R.J., Mate B.R. and Ollervides U.F. (2000) High gray whale mortality and low recruitment in 1999: potential causes and implications. *Journal of Cetacean Research and Management* 2, 85–99.

Le Quéré C., Rödenbeck C., Buitenhuis E.T., Conway T.J., Langenfelds R., Gomez A., Labuschagne C., Ramonet M., Nakazawa T., Metzl N., Gillett N. and Heimann M. (2007) Saturation of the Southern

Ocean CO_2 sink due to recent climate change. *Science Magazine* 316, 1735–1738.

Leaman K.D. and Schott F.A. (1991) Hydrographic structure of the convection regime in the Gulf of Lions: winter 1987. *Journal of Physical Oceanography* 21, 575–598.

Leaper R., Cooke J., Trathan P., Reid K., Rowntree V. and Payne R. (2006) Global climate drives southern right whale (*Eubalaena australis*) population dynamics. *Biology Letters* 2, 289–292.

Learmonth J.A., Macleod C.D., Santos M.B., Pierce J.G., Crick H.Q.P. and Robinson R.A. (2006) Potential effects of climate change on marine mammals. *Oceanography and Marine Biology: an Annual Review* 44, 431–464.

Lejeusne C. (2005) *Habitat fragmenté et métapopulations en milieu marin: structures démographiques et génétiques d'invertébrés cavernicoles dans un contexte de réchauffement climatique*. Thèse de doctorat. COM/Université de la Méditerranée, Marseille, France 221 pp.

Lejeusne C. and Chevaldonné P. (2005) Population structure and life history of *Hemimysis margalefi* (Crustacea: Mysidacea), a 'thermophilic' cave-dwelling species benefiting from the warming of the NW Mediterranean. *Marine Ecology Progress Series* 287, 189–199.

Li L. (2003) *Evolution future du climat en Méditerranée: vers un état de sécheresse accru?* Rapport Quadriennal du CNFGG, pp. 220–223.

Li L., Bozec A., Somot S. and Crépon M. (2006) Modelling regional-scale change of the Mediterranean. *CLIVAR Newsletter Exchanges* 37, 24–27.

Licandro P. and Ibanez F. (2000) Changes of zooplankton communities in the Gulf of Tigullio (Ligurian Sea, western Mediterranean) from 1985 to 1995. Influence of hydroclimatic factors. *Journal of Plankton Research* 22, 2225–2253.

Lionello P. and Sanna A. (2005) Mediterranean wave climate variability and its links with NAO and Indian Monsoon. *Climate Dynamics* 25, 611–623.

Litchman E., Klausmeier C.A., Miller J.R., Schofield O.M. and Falkowski P.G. (2006) Multi-nutrient, multi-group model of present and future oceanic phytoplankton communities. *Biogeosciences* 3, 585–606.

Littaye A., Gannier A., Laran S. and Wilson J.P.F. (2004) The relationship between summer aggregation of fin whales and satellite-derived environmental conditions in the northwestern Mediterranean Sea. *Remote Sensing of Environment* 90, 44–52.

Lloret J., Lleonart J., Solé I. and Fromentin J.M. (2001) Fluctuations of landings and environmental conditions in the north-western Mediterranean Sea. *Fisheries Oceanography* 10, 33–50.

Lloret J., Palomera I., Salat J. and Sole I. (2004) Impact of freshwater input and wind on landings of anchovy (*Engraulis encrasicolus*) and sardine (*Sardina pilchardus*) in shelf waters surrounding the Ebre (Ebro) River delta (north-western Mediterranean). *Fisheries Oceanography* 13, 102–110.

Lockyer C.H. (1986) Body fat condition in Northeast Atlantic fin whales, *Balaenoptera physalus*, and its relationship with reproduction and food resource. *Canadian Journal of Fisheries and Aquatic Sciences* 43, 142–147.

Lopez-Jurado J.L. (2002) Interannual variability in waters of the Balearic Islands. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 33–36 [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Louisy P. and Culioli J.-M. (1999) Synthèse des observations sur l'activité reproductive du mérou brun *Epinephelus marginatus* (Lowe, 1834) en Méditerranée nord-occidentale. *Marine Life* 9, 47–57.

Lusseau D., Williams R., Wilson B., Grellier K., Barton T.R., Hammond P.S. and Thompson P.M. (2004) Parallel influences of climate on the behaviour of Pacific killer whales and Atlantic bottlenose dolphins. *Ecology Letters* 7, 1068–1076.

Luterbacher J., Dietrich D., Xoplaki E., Grosjean M. and Wanner H. (2004) European seasonal and annual temperature variability—trends and extremes since 1500. *Science* 303, 1499–1503.

Luterbacher J., Liniger M.A., Menzel A., Estrella N., Della-Marta P.M., Pfister C., Rutishauser T. and Xoplaki E. (2007) The exceptional European warmth of autumn 2006 and winter 2007: historical context, the underlying dynamics and its phenological impacts. *Geophysical Research Letters* 34, 1–6.

MacGarvin M. and Simmonds M. (1996) Whales and climate change. In Simmonds M.P. and Hutchinson J.D. (eds) *The conservation of whales and dolphins*. Chichester: John Wiley & Sons Ltd, pp. 321–332.

MacLeod C.D., Bannon S.M., Pierce G.J., Schweder C.S., Learmonth J.A., Herman J.S. and Reid R.J. (2005) Climate change and the cetacean community of north-west Scotland. *Biological Conservation* 124, 477–483.

Macquart-Moulin C. and Patriti G. (1996) Accumulation of migratory micronekton crustaceans over the upper slope and submarine canyons of the Northwestern Mediterranean. *Dee-Sea Research* 43, 579–601.

Manca B., Klein B., Kress N. and Ribera D'Alcala M. (2002) Low-frequency changes of water masses structure, flow patterns and biochemical exchanges through the Eastern Mediterranean regions. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 61–70. [Available on line at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Manzanilla S.R. (1989) The 1982–1983 El Niño event recorded in dentinal growth layers in teeth of Peruvian dusky dolphins (*Lagenorhynchus obscurus*). *Canadian Journal of Zoology* 67, 2120–2125.

Marcogliese D.J. (2001) Implications of climate change for parasitism of animals in the aquatic environment. *Canadian Journal of Zoology* 79, 1331–1352.

Mariotti A., Zeng N. and Lau K.M. (2002) Euro-Mediterranean rainfall and ENSO—a seasonally varying relationship. *Geophysical Research Letters*, 29, 591–594.

Maze K.S. and Würsig B. (1999) Bottlenose dolphins of San Luis Pass, Texas: occurrence patterns, site fidelity, and habitat use. *Aquatic Mammals* 25, 91–103.

Meinesz A., Belsher T., Thibaut T., Antolic B., Mustapha K.B., Boudouresque C.F., Chiaverini D., Cinelli F., Cottalorda J.M., Djellouli A., El Abed A., Orestano C., Grau A.M., Ivesa L., Jaklin A., Langar H., Massuti-Pascual E., Peirano A., Tunesi L., de Vaugelas J., Zavodnik N. and Zuljevic A. (2001) The introduced green alga *Caulerpa taxifolia* continues to spread in the Mediterranean. *Biological Invasions* 3, 201–210.

Mercado J.M., Ramirez T., Cortes D., Sebastian M. and Vargas-Yanez M. (2005) Seasonal and inter-annual variability of the phytoplankton communities in an upwelling area of the Alboran Sea (SW Mediterranean Sea). *Scientia Marina* 69, 451–465.

Michaelidis B., Ouzounis C., Paleras A. and Portner H.O. (2005) Effects of long-term moderate hypercapnia on acid–base balance and growth rate in marine mussels *Mytilus galloprovinciallis*. *Marine Ecology Progress Series* 293, 109–118.

Millot C. and Taupier-Letage I. (2004) The cetacean world as seen by physical and biological oceanographers. In *Investigating the role of cetaceans in marine ecosystems, CIESM Workshop Monograph 25,*

SC/N10/CC For Info 2

Monaco, pp. 107–116. [Available at: http://www.ciesm.org/online/monographs/Venice.html; accessed 7 June 2007.]

Millot C., Candela J., Fuda J.L. and Tber Y. (2006) Large warming and salinification of the Mediterranean outflow due to changes in its composition. *Deep-Sea Research* 53, 656–666.

Mills C. (1995) Medusae, siphonophores, and ctenophores as planktivorous predators in changing global ecosystems. *ICES Journal of Marine Science* 52, 575–581.

Moline M.A., Claustre H., Frazer T.K., Schofields O. and Vernet M. (2004) Alterations of the food web along the Antarctic Peninsula in response to a regional warming trend. *Global Change Biology* 10, 1973–1980.

Molinero J.C., Ibanez F., Nival P., Buecher E. and Souissi S. (2005a) North Atlantic climate and Northwestern Mediterranean plankton variability. *Limnology and Oceanography* 50, 164–171.

Molinero J.C., Ibanez F., Souissi S., Chifflet M. and Nival P. (2005b) Phenological changes in the Northwestern Mediterranean copepods *Centropages typicus* and *Temora stylifera* linked to climate forcing. *Oecologia* 145, 640–649.

Molinero J.C., Ibanez F., Souissi S., Licandro P., Buecher E., Dallot S. and Nival P. (2007) Northern Hemisphere climate impact on Mediterranean zooplankton. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 7 June 2007.]

Monaco A. (1998) *La Méditerranée, zone test des changements globaux et régionaux des ecosystems. Lettre No. 8 du Programme International Géosphère Biosphère-Programme Mondial de Recherches sur le Climat (PIGB-PMRC).* 8 p. [Available at: http://www.cnrs.fr/cw/dossiers/dosclim/rechfran/4theme/roledeLocean/mediterranee.html; accessed 4 June 2007.]

Moore S.E., Grebmeier J.M. and Davies J.R. (2003) Gray whale distribution relative to forage habitat in the northern Bering Sea: current conditions and retrospective summary. *Canadian Journal of Zoology* 81, 734–742.

Morri C. and Bianchi C.N. (2001) Recent changes in biodiversity in the Ligurian Sea (NW Mediterranean): is there a climatic forcing? In Faranda F.M., Guglielmo L. and Spezie G. (eds) *Structure and processes in the Mediterranean ecosystems*. Milan: Springer, pp. 375–384.

Morton A.B. (2000) Occurrence, photo-identification and prey of Pacific white-sided dolphins (*Lagenorhynchus obliquidens*) in the Broughton Archipelago, Canada 1985–1997. *Marine Mammal Science* 16, 80–93.

Nehring S. (1998) Establishment of thermophilic phytoplankton species in the North Sea: biological indicators of climatic changes. *ICES Journal of Marine Science* 55, 818–823.

Neumann D.R. (2001) Seasonal movements of short-beaked common dolphins (*Delphinus delphis*) in the north-western Bay of Plenty, New Zealand: influence of sea surface temperature and El Niño and La Niña. *New Zealand Journal of Marine and Freshwater Research* 35, 371–374.

Nicholls R.J. and Hoozemans F.M.J. (1996) The Mediterranean: vulnerability to coastal implications of climate change. *Ocean and Coastal Management* 31, 105–132.

Northridge S. (1984) *World review of interactions between marine mammals and fisheries*. FAO Fisheries Technical Paper 251. Rome: Food and Agriculture Organization. 190 pp.

Notarbartolo di Sciara G., Zanardelli M., Jahoda M., Panigada S. and Airoldi S. (2003) The fin whale *Balaenoptera physalus* (L. 1758) in the Mediterranean Sea. *Mammal Review* 33, 105–150.

Occhipinti-Ambrogi A. (2007) Global change and marine communities: alien species and climate change. *Marine Pollution Bulletin* 55, 342–352.

Ohman M.D. and Hirche H.J. (2001) Density-dependent mortality in an oceanic copepod population. *Nature* 412, 638–641.

Olita A., Sorgente R., Natale S., Gabersek S., Ribotti A., Bonanno A. and Patti B. (2007) Effects of the 2003 European heatwave on the central Mediterranean Sea: surface fluxes and the dynamical response. *Ocean Science* 3, 273–289.

Oren O.H. (1957) Changes in the temperature of the Eastern Mediterranean Sea in relation to the catch of the Israel trawl fishery during the years 1954/55 and 1955/56. *Bulletin de l'Institut Océanographique (Monaco)* 1102, 1–12.

Orr J.C., Maier-Reimer E., Mikolajewicz U., Monfray P., Ray G.C., Hayden B.P., Bulger A.J. and McCormick-Ray M.G. (1992) Effects of global warming on the biodiversity of coastal-marine zones. In. Peters R.L. and Lovejoy T.E. (eds) *Global warming and biological diversity*. New Haven, CT: Yale University Press, pp. 91–105.

Orr J.C., Maier-Reimer E., Mikolajewicz U., Monfray P., Sarmiento J.L., Toggweiler J.R., Taylor N.K., Palmer J., Gruber N., Sabine C.L., Le Quere C.L., Key R.M. and Boutin J. (2001) Estimates of anthropogenic carbon uptake from four three-dimensional global ocean models, *Global Biogeochemical Cycles* 15, 43–60.

Orr J.C., Fabry V.J., Aumont O., Bopp L., Doney S.C., Feely R.A., Gnanadesikan A., Gruber N., Ishida A., Joos F., Key R.M., Lindsay K., Maier-Reimer E., Matear R., Monfray P., Mouchet A., Najjar R.G., Plattner G.K., Rodgers K.B., Sabine C.L., Sarmiento J.L., Schlitzer R., Slater R.D., Totterdell I.J., Weirig M.F., Yamanaka Y. and Yool A. (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437, 681–686.

Orsi Relini L. and Giordano M. (1992) The fin whale and other large pelagic filters as samplers of *Meganyctiphanes norvegica*. *Rapport de la CIESM* 33, 263.

Orsi Relini L., Relini G., Cima C., Palandri G., Relini M. and Torchia G. (1994) *Meganyctiphanes norvegica* and fin whales in the Ligurian Sea: new seasonal patterns. *European Research on Cetaceans* 8, 179–182.

Orsi Relini L. and Relini M. (1993) The stomach content of some common dolphins (*Delphinus delphis* L.) from the Ligurian Sea. *European Research on Cetaceans* 7, 99–102.

Oviatt C.A. (2004) The changing ecology of temperate coastal waters during a warming trend. *Estuaries* 27, 895–904.

Palka D., Read A. and Potter C. (1997) Summary of knowledge of white-sided dolphins (*Lagenorhynchus acutus*) from US and Canadian Atlantic waters. *Report of the International Whaling Commission* 47, 729–734.

Parenti P. and Bressi N. (2001) First record of the orange spotted grouper, *Epinephelus coioides* (Perciformes: Serranidae) in the northern Adriatic Sea. *Cybium* 25, 281–284.

Parmesan C. and Yohe G. (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421, 37–42.

Parry M.L. (ed.) (2000) *Assessment of potential effects and adaptation for climate change in Europe: The Europe Acacia Project*. Jackson Environmental Institute, University of East Anglia, Norwich, UK.

Paskoff R. (2001) *L'élévation du niveau de la mer et les espaces côtiers*. Paris: Propos-Institut Océanographique.

Pechenik J.A. (1989) Environmental influences on larval survival and development. In Giese A.C., Pearse J.S. and Pearse V.B. (eds) *Reproduction of marine invertebrates*. Palo Alto, CA: Blackwell Scientific Publications, pp. 551–608.

Peirano A., Damasso V., Montefalcone M., Morri C. and Bianchi C.N. (2005) Effects of climate, invasive species and anthropogenic impacts on the growth of seagrass *Posidonia oceanica* (L) Delile in Liguria (NW Mediterranean Sea). *Marine Pollution Bulletin* 50, 817–822.

Perez T., Garrabou J., Sartoretto S., Harmelin J.G., Francour P. and Vacelet J. (2000) Mortalité massive d'invertébrés marins: unévénement sans précédent en Méditerranée nord-occidentale. *Comptes Rendus de l' Académie des Sciences Paris, Sciences et Vie* 323, 853–865.

Perrin W.F. (1989) *Dolphins, porpoises, and whales—an action plan for the conservation of biological diversity: 1988–1992*. Second edition. IUCN/SSC Cetacean Specialist Group. 30 pp.

Perry A.L., Low P.J., Ellis J.R. and Reynolds J.D. (2005) Climate change and distribution shifts in marine fishes. *Science* 308, 1912–1924.

Petchey O.L., McPhearson P.T., Casey T.M. and Morin P.J. (1999) Environmental warming alters food-web structure and ecosystem function. *Nature* 402, 69–72.

Pfeifle G., Mays C. and Poumadère M. (2004) Changement climatique, augmentation du niveau de la mer et (ré)action des parties prenantes: étude de cas à l'aide d'un scénario catastrophe en Camargue. In Scarwell H.J. and Franchomme M. (eds) *Contraintes environnementales et gouvernance des territoires*. Lille: Editions de l'Aube (Nord).

Philippart C.J.M., Van Aken H.M., Beukema J.J., Bos O.G., Cadee G.C. and Dekker R. (2003) Climate-related changes in recruitment of the bivalve *Macoma balthica*. *Limnology and Oceanography* 48, 2171–2185.

Pierce G.J. and Boyle P.R. (2003) Empirical modeling of inter-annual trends in abundance of squid (*Loligo forbesi*) in Scottish waters. *Fisheries Research* 59, 305–326.

Piontkovski S.A. and Castellani C. (2007) Decline of zooplankton biomass in the Tropical Atlantic Ocean. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 7 June 2007.]

Pitois S.G. and Fox C.J. (2007) Have climate induced changes in zooplankton community led to poor conditions for larval fish growth? A test on cod larvae on the UK shelf. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 7 June 2007.]

Planque B. and Taylor A.H. (1998) Long-term changes in zooplankton and the climate of the North Atlantic. *ICES Journal of Marine Science* 55, 644–654.

Polat S. (2004) New record for a dinoflagellate species (*Citharistes regius* Stein) in the Northern Levantine Basin (Eastern Mediterranean). *Turkish Journal of Botany* 28, 507–509.

Politi E. (1998) Un progetto per i delfini in Mediterraneo. *Le Scienze* 360, 64–69.

Politi E., Bearzi G. and Airoldi S. (2000) Evidence for malnutrition in bottlenose dolphins photoidentified in the eastern Ionian Sea. *European Research on Cetaceans* 14, 234–236.

Politi E. and Bearzi G. (2004) Evidence of rarefaction for coastal common dolphins in the eastern Ionian Sea. *European Research on Cetaceans* 15, 449–452.

Pörtner H.O. and Knust R. (2007) Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science* 315, 95–97.

Pörtner H.O., Langenbuch M. and Reipschläger A. (2004) Biological impact of elevated ocean CO_2 concentrations: lessons from animal physiology and earth history. *Journal of Oceanography* 60, 705–718.

Poulard J.C. and Blanchard F. (2005) The impact of climate change on the fish community structure of the eastern continental shelf of the Bay of Biscay. *ICES Journal of Marine Science* 62, 1436–1443.

Poumadère M., Mays C., Le Mer S. and Blong R. (2005) Dangerous climate here and now: the 2003 heat wave in France. *Risk Analysis* 25, 1483–1494.

Pozo-Vasquez D., Esteban-Parra M.J., Rodrigo F.S. and Castro-Diez Y. (2001) A study of NAO variability and its possible non-linear influences on European surface temperature. *Climate Dynamics* 17, 701–715.

Prieur L. (2002) Physical historical data on the Ligurian Sea from the Villefranche-sur-Mer Observatory. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 37–40. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Pustelnik G. (1976) *Les Euphausiacés dans l 'écosystème épipélagique de la Mer Ligure. Méthodologie. Etude* in situ. Mémoire thèse de fin d'étude. Centre de Recherches Océanographique de Villefranche sur Mer. Rapport 18, 62 pp.

Quynh V.D. (1978) *Ecologie des crustacés micronectoniques dans l'écosystème superficiel du bassin liguro-provençal: étude particulière des décapodes Natantia pélagiques*. Thèse de 3° cycle Université Pierre et Marie (Paris 6), 125 pp.

Raicich F., Pinardi N. and Navarra A. (2001) Teleconnections between Indian monsoon and Sahel rainfall and the Mediterranean. *Archivio di Oceanografia e Limnologia* 22, 9–14.

Raniello R., Lorenti M., Brunet C. and Buia M.C. (2004) Photosynthetic plasticity of an invasive variety of *Caulerpa racemosa*: light harvesting capacity and seasonal acclimation. *Marine Ecology Progress Series* 271, 113–120.

Reeves R.R. (1991) Speculations on the impact of global warming on aquatic mammals. In Taylor S. (ed.) *They're not saved yet. Proceedings of the American Cetacean Society, fourth biennial conference? Monterey, California, November 9–11, 1990*. San Pedro, California: American Cetacean Society, pp. 19–23.

Read A. and Gaskin D.E. (1990) Changes in growth and reproduction of harbour porpoises, *Phocoena phocoena*, from the Bay of Fundy. *Canadian Journal of Fisheries and Aquatic Sciences* 47, 2158–2163.

Reeves R.R. and Notarbartolo di Sciara G. (2006) *The status and distribution of cetaceans in the Black Sea and Mediterranean Sea*. Workshop Report, 5–7 March, Monaco. 142 pp.

Reeves R.R., Smith B.D., Crespo E. and Notarbartolo di Sciara G. (2003) *Dolphins, whales, and porpoises: 2002–2010 Conservation Action Plan for the world's cetaceans*. Gland, Switzerland: UICN. 139 pp.

Regner S. (1996) Effects of environmental changes on early stages and reproduction of anchovy in the Adriatic Sea. *Scientia Marina* 60 (Supplement 2), 167–177.

Relini M. and Orsi Relini L. (1997) The two species of barracuda (Sphyraenidae) in the western Mediterranean. *Cybium* 21, 216–222.

SC/N10/CC For Info 2

Ricciardi A. (2004) Assessing species invasions as a cause of extinction. *Trends in Ecology and Evolution* 19, 619.

Riebesell U., Zondervan I., Rost B., Tortell P.D., Zeebe R.E. and Morel F.M.M. (2000) Reduced calcification of marine plankton in response to increased atmospheric CO_2. *Nature* 407, 364–367.

Rixen M., Beckers J.M., Levitus S., Antonov J., Boyer T., Maillard C., Fichaut M., Balopoulos E., Iona S., Dooley H., Garcia M.J., Manca B., Giorgetti A., Manzella G., Mikhailov N., Pinardi N. and Zavatatrelli M. (2005) The Western Mediterranean deep water: a proxy for climate change. *Geophysical Research Letters* 32, L12608, doi:10.1029/2005GL022702.

Robinson R.A., Learmouth J.A., Hutson A.M., Macleod C.D., Sparks T.H., Leech D.I., Pierce G.J., Rehfisch M.M. and Crick H.Q.P. (2005) *Climate change and migratory species. BTO Research Report 414.* Thetford, UK: British Trust for Ornithology. [Available at: http://www.defra.gov.uk/wildlifecountryside/resprog/findings/climatechange-migratory/climatechange-migratory.pdf, accessed 4 March 2007.]

Roemmich D. and McGowan J.A. (1995) Climatic warming and the decline of zooplankton in the California current. *Science* 267, 1324–1326.

Roessig J., Woodley C., Cech J. and Hansen L. (2004) Effects of global climate change on marine and estuarine fishes and fisheries. *Reviews in Fish Biology and Fisheries* 14, 251–275.

Roether W., Manca B., Klein B., Bregant D., Georgopoulos D., Beitzel V., Kovacevic V. and Luchetta A. (1996) Recent changes in Eastern Mediterranean deep waters. *Science* 271, 333–335.

Rohling E.J. and Bryden H.L. (1992) Man-induced salinity and temperature increases in the Western Mediterranean deep water. *Journal of Geophysical Research* 97, 11191–11198.

Romano J.C., Bensoussan N., Younes W. and Arlhac D. (2000) Anomalie thermique dans les eaux du golfe de Marseille durand l'été 1999. Une explication partielle de la mortalite d'invertébrés fixés? *Comptes Rendud de l' Académie des Sciences Paris, Sciences et Vie* 323, 415–427.

Root T.L., Price J.T., Hall K.R., Schneider S.H., Rosenzweig C. and Pounds J.A. (2003) Fingerprints of global warming on wild animals and plants. *Nature* 421, 57–60.

Rose G.A. (2005) On distributional responses of North Atlantic fish to climate change. *ICES Journal of Marine Science* 62, 1360–1374.

Ross P.S., Vos J.G., Birnbaum L.S. and Osterhaus A.D.M.E. (2000) PCBs are a health risk for humans and wildlife. *Science* 289, 1878–1879.

Royal Society (2005) *Ocean acidification due to increasing atmospheric carbon dioxide. Policy Document 12/05*. London: The Royal Society, 60 pp.

Ruitton S., Javel F., Culioli J.-M., Meinesz A., Pergent G. and Verlaque M. (2005) First assessment of the *Caulerpa racemosa* (Caulerpales, Chlorophyta) invasion along the French Mediterranean coast. *Marine Pollution Bulletin* 50, 1061–1068.

Russo A., Rabitti S. and Bastianini M. (2002) Decadal climatic anomalies in the northern Adriatic Sea inferred from a new oceanographic data set. *PSZNI Marine Ecology* 23, 340–351.

Sabates A. and Gili J.M. (1991) Processes occurring during fish larvae stages as determinants of year-class strength in fish population in the northwestern Mediterranean. *Oceanologica Aquatica* 10, 203–213.

Sabates A., Martin P., Loret J. and Raya V. (2006) Sea warming and fish distribution: the case of the small pelagic fish, *Sardinella aurita*, in the western Mediterranean. *Global Change Biology* 12, 2209–2219.

Salat J. (1996) Review of hydrographic environmental factors that may influence anchovy habitats in northwestern Mediterranean. *Scientia Marina* 60, 21–32.

Salat J. and Pascual J. (2002) The oceanographic and meteorological station at L'Estartit (NW Mediterranean). In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 29–32. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Sanchez P. (1982) Régimen alimentario de *Illex coindetii* (Verany, 1837) en el mar Catalan. *Investigacion Pesquera* 46, 443–449.

Sanford E. (1999) Regulation of keystone predation by small changes in ocean temperature. *Science* 283, 2095–2097.

Scarnecchia D.L. (1984) Climate and oceanic variations affecting yield of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* 41, 917–935.

Schiel D.R., Steinbeck J.R. and Foster M.S. (2004) Ten years of induced ocean warming causes comprehensive changes in marine benthic communities. *Ecology* 85, 1833–1839.

Sebens K.P. (1994) Biodiversity of coral reefs: what are we loosing and why? *American Zoologist* 34, 115–133.

Serena F. and Silvestri R. (1996) First record of *Pomadasys incisus* (Haemulidae) in the Northern Tyrrhenian Sea. *Cybium* 20, 409–411.

Shane S.H. (1990) Comparison of bottlenose dolphin behavior in Texas and Florida, with a critique of methods for studying dolphin behavior. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. San Diego, CA: Academic Press, pp. 541–558.

Shane S.H. (1995) Relationship between pilot whales and Risso's dolphins at Santa Catalina Island, California, USA. *Marine Ecology Progress Series* 123, 5–11.

Sheppard C.R.C. (1999) Coral decline and weather patterns over 20 years in the Chagos Archipelago, Central Indian Ocean. *Ambio* 28, 472–478.

Shirayama Y. and Thornton H. (2005) Effect of increased atmospheric CO_2 on shallow water marine benthos. *Journal of Geophysical Research* 110, C09S08, doi:10.1029/2004JC002618.

Silva M.A. (1999) Diet of common dolphins, *Delphinus delphis*, off the Portuguese coast. *Journal of the Marine Biological Association of the United Kingdom* 79, 531–540.

Simmonds M.P. (1994) Saving Europe's dolphins. *Oryx* 28, 238–248.

Simmonds M.P. (2004) *Whales and dolphins of the world.* London: New Holland Publishers.

Simmonds M.P. and Nunny L. (2002) Cetacean habitat loss and degradation in the Mediterranean Sea. In Notarbartolo di Sciara G. (ed.) *Cetaceans of the Mediterranean and Black Seas: state of knowledge and conservation strategies*. A report to the ACCOBAMS Secretariat, Monaco, February 2002. Section 7, 23 pp.

Simmonds M.P. and Mayer S.J. (1997) An evaluation of environmental and other factors in some recent marine mammal mortalities in Europe: implication for conservation and management. *Environmental Reviews* 5, 89–98.

Simmonds M.P. and Isaac S.J. (2007) The impacts of climate change on marine mammals: early signs of significant problems. *Oryx* 41, 19–26.

Sims D.W. and Reid P.C. (2002) Congruent trends in long-term zooplankton decline in the north-east Atlantic and basking shark (*Cetorhinus maximus*) fishery catches off west Ireland. *Fisheries Oceanography* 11, 59–63.

Sims D.W., Genner M.J., Southward A.J. and Hawkins S.J. (2001) Timing of squid migration reflects North Atlantic climate variability. *Proceedings of the Royal Society of London Series B: Biological Sciences* 268, 2607–2611.

SC/N10/CC For Info 2

Sims D.W., Wearmouth V.J., Genner M.J., Southward A.J. and Hawkins S.J. (2004) Low-temperature-driven early spawning migration of a temperate marine fish. *Journal of Animal Ecology* 73, 333–341.

Sissener E.H. and Bjorndal T. (2005) Climate change and the migratory pattern for Norwegian spring-spawning herring—implications for management. *Marine Policy* 29, 299–309.

Slagstad D., Ellingsen I. and Edvardsen A. (2007) Effect of climatic change on the production and distribution of *C. finmarchicus* in the Norwegian and Barents Seas. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 7 June 2007.]

Smith S.C. and Whitehead H. (1993) Variations in the feeding success and behaviour of Galapagos sperm whale (*Physeter macrocephalus*) as they relate to oceanographic conditions. *Canadian Journal of Zoology* 71, 1991–1996.

Smith L.C., Sheng Y., MacDonald G.M. and Hinzman L.D. (2005) Disappearing Arctic lakes. *Science* 308, 1429.

Somero G.N. (2002) Thermal physiology and vertical zonation of intertidal animals: optima, limits, and costs of living. *Integrative and Comparative Biology* 42, 780–789.

Somot S., Sevault F. and Déqué M. (2006) Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model. *Climate Dynamics* 27, 851–879.

Southward A.J., Boalch G.T. and Mattock L. (1988) Fluctuations in the herring and pilchard fisheries of Devon and Cornwall linked to change in climate since the 16th century. *Journal of the Marine Biological Association of the United Kingdom* 68, 423–445.

Southward A.J., Hawkins S.J. and Burrows M.T. (1995) Seventy years' observations of changes in distribution and abundance of zooplankton and intertidal organisms in the western English Channel in relation to rising sea temperature. *Journal of Thermal Biology* 20, 127–155.

Spalding M.D., Teleki K. and Spencer T. (2003) Climate change and coral bleaching. In Green R.E., Harley M., Spalding M.D. and Zöckler C. (eds) *Impacts of climate change on wildlife*. Sandy, UK: RSPB, pp. 38–39.

Stachowicz J.J., Fried H., Osman R.W. and Whitlatch R.B. (2002a) Biodiversity, invasion resistance, and marine ecosystem function: reconciling pattern and process. *Ecology* 83, 2575–2590.

Stachowicz J.J., Terwin J.R., Whitlatch R.B. and Osman R.W. (2002b) Linking climate change and biological invasions: ocean warming facilitates nonindigenous species invasions. *Proceedings of the National Academy of Sciences of the USA* 99, 15497–15500.

Stanley S.M. (1984) Marine mass extinctions: a dominant role for temperature. In Nitecki M.H. (ed.) *Extinctions*. Chicago: University of Chicago Press, pp. 69–117.

Stenevik E.K. and Sundby S. (2007) Impacts of climate change on commercial fish stocks in Norwegian waters. *Marine Policy* 31, 19–31.

Stenseth N., Mysterud A., Ottersen G., Hurrell J.W., Chan K.S. and Lima M. (2002) Ecological effects of climate fluctuations. *Science* 297, 1292–1296.

Stern S.J. (1996) Potential impacts of global change on foraging dynamics and distribution of baleen whales off northern and central California. *The International Whaling Commission's 1996 Annual Meeting, June 23–28, Aberdeen, Scotland*.

Streftaris N. and Zenetos A. (2006) Alien marine species in the Mediterranean—the 100 'worst invasives' and their impact. *Mediterranean Marine Science* 7, 87–118.

Struglia M.V., Mariotti A. and Filograsso A. (2004) River discharge into the Mediterranean Sea: climatology and aspects of the observed variability. *Journal of Climate* 17, 4740–4751.

Svendsen E. (1995) Influence of climate on recruitment and migration of fish stocks in the North Sea. *Canadian Special Publication of Fisheries and Aquatic Sciences* 121, 641–653.

Taleb H., Vale P., Jaime E. and Blaghen M. (2001) Study of paralytic shellfish poisoning toxin profile in shellfish from the Mediterranean shore of Morocco. *Toxicon* 39, 1855–1861.

Tett S.F.B., Jones G.S., Stott P.A., Hill D.C., Mitchell J.F.B., Allen M.R., Ingram W.J., Johns T.C., Johnson C.E., Jones A., Roberts D.L., Sexton D.M.H. and Woodage M.J. (2000) Estimation of natural and anthropogenic contributions to 20th century temperature change. *Journal of Geophysical Research* 107(D16).

Theocharis A., Nittis K., Kontoyiannis H., Papageorgiou E. and Balopoulos E. (1999) Climatic changes in the Aegean Sea influence the thermohaline circulation of the Eastern Mediterranean (1986–1997). *Geophysical Research Letters* 26, 1617–1620.

Thomas D., Cameron A., Green R.E., Bakkenes M., Beaumont L.J., Collingham Y.C., Erasmus B.F.N., Ferreira de Siqueira M., Grainger A., Hannah L., Hughes L., Huntley B., Van Jaarsveld A.S., Midgley G.F., Miles L., Ortega-Huerta M.A., Peterson A.T., Phillips O.L. and Williams S.E. (2004) Extinction risk from climate change. *Nature* 427, 145–148.

Timmermann A., Oberhuber J., Bacher A., Esch M., Latif M. and Roeckner E. (1999) Increased El Niño frequency in a climate model forced by future greenhouse warming. *Nature* 398, 694–697.

Tsimplis M., Zervakis V., Josey S., Peneva E., Struglia M.V., Stanev E., Lionello P., Artale V., Theocharis A., Tragou E. and Rennell J. (2006) Changes in the oceanography of the Mediterranean Sea and their link to climate variability. In Lionello P., Malanotte-Rizzoli P. and Boscolo R. (eds) *Mediterranean climate variability*. Amsterdam: Elsevier, pp. 227–282.

Tsimplis M.N. and Rixen M. (2002) Sea level in the Mediterranean Sea: the contribution of temperature and salinity changes. *Geophysical Research Letters* 29, 51.1–51.4.

Tsimplis M.N. and Josey S.A. (2001) Forcing of the Mediterranean Sea by atmospheric oscillations over the North Atlantic. *Geophysical Research Letters* 28, 803–806.

Turley C.M. (1999) The changing Mediterranean Sea—a sensitive ecosystem? *Progress in Oceanography* 44, 387–400.

UNEP (1996) *The state of the marine and coastal environment in the Mediterranean region. MAP Technical Report Series No. 100*. Athens: UNEP.

UNEP/IUCN (1994) *Technical report on the state of cetaceans in the Mediterranean. Mediterranean Action Plan Technical Reports Series No. 82*. Tunis: UNEP, Regional Activity Centre for Specially Protected Areas. 37 pp.

Vacchi M. and Cau A. (1986) The occurrence of *Sphoeroides cutaneus* (Gunther, 1870) (Pisces, Tetraodontidae) in the middle-west Mediterranean Sea. *Cybium* 10, 199–203.

Vacchi M., Morri C., Modena M., La Mesa G. and Bianchi C. N. (2001) Temperature changes and warm-water species in the Ligurian Sea: the case of the ornate wrasse *Thalassoma pavo* (Linnaeus, 1758). *Archivio di Oceanografia e Limnologia* 22, 149–154.

Valiela I. (1995) *Marine ecological processes*. 2nd edition. New York: Springer. 686 pp.

Van de Bildt M.W.G., Vedder E.J., Martina B.E.E., Abou-Sidib B., Jiddou A.B., Barham M.E.O., Androukaki E., Komnenou A., Niesters H.G.M. and Osterhaus A.D.M.E. (1999) Morbilliviruses in Mediterranean monk seals. *Veterinary Microbiology* 69, 19–21.

SC/N10/CC For Info 2

Vargas-Yanes M., Ramirez T., Cortes D., De Puelles M.L.F., Lavin A., Lopez-Jurado J.L, Gonzales-Pola C., Vidal I. and Sebastian M. (2002) Variability of the Mediterranean water around the Spanish coast: project RADIALES. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 25–28 [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Velsch J.P. (1997) *Dynamique de population et migrations verticales journalières de* Meganyctiphanes norvegica *(Crustacés: Euphausiaces) en Méditerranée nord-occidentale*. Thèse de doctorat. Université de la Méditerranée Aix Marseille, Marseille, France. 129 pp.

Viale D. (1985) Cetaceans in the Northwestern Mediterranean: their place in the ecosystem. *Oceanography and Marine Biology: an Annual Review* 23, 491–571.

Vila M., Garcés E., Maso M. and Camp J. (2001) Is the distribution of the toxic dinoflagellate *Alexandrium catenella* expanding along the NW Mediterranean coast? *Marine Ecology Progress Series* 222, 73–83.

Vilibic I. (2002) Long-term changes in Adriatic thermohaline properties. In *Tracking long-term hydrological change in the Mediterranean Sea. CIESM Workshop Series, 16, 22–24 April, Monaco*, pp. 57–59. [Available at: http://www.ciesm.org/publications/Monaco02.pdf; accessed 7 June 2007.]

Visbeck M.H., Hurrell J.W., Polvani L. and Cullen H.M. (2001) The North Atlantic Oscillation: past, present, and future. *PNAS* 98, 12876–12877.

Voarino L. (2006) *Analyse temporelle du zooplancton méditerranéen dans la rade de Villefranche-sur-Mer. Influence des facteurs environnementaux*. Mémoire de Master, Laboratoire Océanologique de Villefranche-sur-Mer (LOV), Villefranche-sur-Mer, France. [Available at: http://www.obs-vlfr.fr/LOV/ZooPart/ZooScan/Documents_Zooscan/Rapports/rapport_Luna.pdf; accessed 5 May 2007.]

Voss R., Schmidt J.O. and Schnack D. (2007) Impacts of climate change on the Baltic zooplankton–fish larvae link: evidence for sprat (*Sprattus sprattus* L.) larval behaviour change influencing recruitment levels. In *4th International Zooplankton Symposium: human and climate forcing of zooplankton population, Hiroshima, 28 May–1 June 2007. Temporal and regional responses of zooplankton to global warming: phenology and poleward displacement*, pp. 207–212. [Available at: http://www.pices.int/publications/book_of_abstracts/2007_4th_Zooplankton_Symposium.pdf; accessed 7 June 2007.]

Walther G.R., Post E., Convey P., Menzel A., Parmesan C., Beebe T.J.C., Fromentin J.M., Hoegh-Guldberg O. and Bairlein F. (2002) Ecological responses to recent climate change. *Nature* 416, 389–395.

Waluda C.M., Rodhouse P.G., Podestá G.P., Trathan P.N. and Pierce G.J. (2001) Surface oceanography of the inferred hatching grounds of *Illex argentinus* (Cephalopoda: Ommastrephidae) and influences on recruitment variability. *Marine Biology* 139, 671–679.

Wells R.S., Hansen L.J., Baldridge A., Dohl T.P., Kelly D.L. and Defran R.H. (1990) Northward extension of the range of bottlenose dolphins along the California coast. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. San Diego: Academic Press, pp. 421–431.

Whitehead H. (1997) Sea surface temperature and the abundance of sperm whale calves off the Galápagos Islands: implications for the effects of global warming. *Report of the International Whaling Commission* 47, 941–944.

Wiley D.N. and Clapham P.J. (1993) Does maternal condition affect the sex ratio of offspring in humpback whales? *Animal Behaviour* 46, 321–324.

Wilson B., Arnold H., Bearzi G., Fortuna C.M., Gaspar R., Ingram S., Liret C., Pribanic S., Read A.J., Ridoux V., Schneider K., Urian K.W., Wells R.S., Wood C., Thompson P.M. and Hammond P.S. (1999) Epidermal diseases in bottlenose dolphins: impacts of natural and anthropogenic factors. *Proceedings of the Royal Society of London Series B, Biological Sciences* 266, 1077–1083.

Wilson B., Reid R.J., Grellier K., Thompson P.M. and Hammond P.S. (2004) Considering the temporal when managing the spatial: a population range expansion impacts protected areas-based management for bottlenose dolphins. *Animal Conservation* 7, 331–338.

Woodwell G. and Mackenzie F. (1995) *Biotic feedbacks in the global climate system: will the warming feed the warming?* New York: Oxford University Press.

Worms B., Sandow M., Oschlies A., Lotze H.K. and Myers R.A. (2005) Global patterns of predator diversity in the open oceans. *Science* 309, 1365–1369.

Würsig B., Reeves R.R. and Ortega-Ortiz J.G. (2002) Global climate change and marine mammals. In Evans P.G.H. and Raga J.A. (eds) *Marine mammals: biology and conservation*. New York: Kluwer Academic/Plenum Publishers, pp. 589–608.

Wurtz M. and Marrale D. (1991) On the stomach contents of striped dolphins (*Stenella coeruleoalba*, Meyen 1838) from the Ligurian coast, central Mediterranean Sea. In *Proceedings of the Fifth Annual Conference of the European Cetacean Society, Sandefjord, Norway, 21–23 February 1991*, pp. 62–64.

Xoplaki E., Gonzalez-Rouco J.F., Luterbacher J. and Wanner H. (2004) Wet season Mediterranean precipitation variability: influence of large-scale dynamics and trends. *Climate Dynamics* 23, 63–78.

Xoplaki E., Luterbacher J., Paeth H., Dietrich D., Steiner N., Grosjean M. and Wanner H. (2005) European spring and autumn temperature variability and change of extremes over the last half millennium. *Geophysical Research Letters* 32, 1–2.

Xoplaki E., Luterbacher J. and Gonzalez-Rouco F.J. (2006) Mediterranean summer temperature and winter precipitation, large-scale dynamics, trends. *Il Nuovo Cimento C* 29, 45–54.

Zabala M., Garcia-Rubies A., Louisy P. and Sala A. (1997) Spawning behaviour of the Mediterranean dusky grouper *Epinephelus marginatus* (Lowe, 1934) (Pisces, Serranidae) in the Medes Islands Reserve (NW Mediterranean, Spain). *Scientia Marina* 61, 65–77.

Zaitsey Y. and Mamaev V. (1997) *Marine biological diversity in the Black Sea: a study of change and decline*. New York: United Nations Publications.

Zenetos A., Siokou-Frangou I., Gotsis-Skretas O. and Groom S. (2002) *Europe's biodiversity–biogeographical regions and seas. The Mediterranean Sea—blue oxygen-rich nutrient-poor waters.* [Available at: http://reports.eea.eu.int/report_2002_0524_154909/en; Accessed 7 June 2007.]

Zenetos A., Todorova V. and Alexandrov B. (2003) Marine biodiversity changes in the Mediterranean and Black Sea regions. *International Conference on the Sustainable Development of the Mediterranean and Black Sea Environment*. Workshop, 28 May–1 June, Thessaloniki, Greece.

Zore-Armanda M. (1991) Natural characteristics and climatic changes of the Adriatic Sea. *Acta Adriatica* 32, 567–586.

and

Zore-Armanda M., Stojanoski L. and Vukadin I. (1987) Time series of oceanographic parameters: eutrophication of the open Adriatic Sea. *FAO Fisheries Report* 394, 71–77.

Correspondence should be addressed to:
D.D. Gambaiani
Souffleurs d'Ecume, Domaine de l'Escarelle
83170 La Celle, France
email: delphine.gambaiani@souffleursdecume.com